

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/mycres

A higher-level phylogenetic classification of the Fungi

David S. HIBBETT^{a,*}, Manfred BINDER^a, Joseph F. BISCHOFF^b, Meredith BLACKWELL^c, Paul F. CANNON^d, Ove E. ERIKSSON^e, Sabine HUHDORF^f, Timothy JAMES^g, Paul M. KIRK^d, Robert LÜCKING^f, H. THORSTEN LUMBSCH^f, François LUTZONI^g, P. Brandon MATHENY^a, David J. MCLAUGHLIN^h, Martha J. POWELLⁱ, Scott REDHEAD^j, Conrad L. SCHOCH^k, Joseph W. SPATAFORA^k, Joost A. STALPERS^l, Rytas VILGALYS^g, M. Catherine AIME^m, André APTROOTⁿ, Robert BAUER^o, Dominik BEGEROW^p, Gerald L. BENNY^q, Lisa A. CASTLEBURY^m, Pedro W. CROUS^l, Yu-Cheng DAI^r, Walter GAMS^l, David M. GEISER^s, Gareth W. GRIFFITH^t, Cécile GUEIDAN^g, David L. HAWKSWORTH^u, Geir HESTMARK^v, Kentaro HOSAKA^w, Richard A. HUMBER^x, Kevin D. HYDE^y, Joseph E. IRONSIDE^t, Urmaz KÖLJALG^z, Cletus P. KURTZMAN^{aa}, Karl-Henrik LARSSON^{ab}, Robert LICHTWARDT^{ac}, Joyce LONGCORE^{ad}, Jolanta MIĄDLIKOWSKA^g, Andrew MILLER^{ae}, Jean-Marc MONCALVO^{af}, Sharon MOZLEY-STANDRIDGE^{ag}, Franz OBERWINKLER^o, Erast PARMASTO^{ah}, Valérie REEB^g, Jack D. ROGERS^{ai}, Claude ROUX^{aj}, Leif RYVARDEN^{ak}, José Paulo SAMPAIO^{al}, Arthur SCHÜßLER^{am}, Junta SUGIYAMA^{an}, R. Greg THORN^{ao}, Leif TIBELL^{ap}, Wendy A. UNTEREINER^{aq}, Christopher WALKER^{ar}, Zheng WANG^a, Alex WEIR^{as}, Michael WEISS^o, Merlin M. WHITE^{at}, Katarina WINKA^e, Yi-Jian YAO^{au}, Ning ZHANG^{av}

^aBiology Department, Clark University, Worcester, MA 01610, USA

^bNational Library of Medicine, National Center for Biotechnology Information, 45 Center Drive, Bethesda, MD 20892-6510, USA

^cDepartment of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

^dCABI UK, Bakeham Lane, Egham, Surrey TW20 9TY, UK

^eDepartment of Ecology and Environmental Science, Umeå University, SE-901 87 Umeå, Sweden

^fDepartment of Botany, The Field Museum of Natural History, 1400 South Lake Shore Drive, Chicago, IL 60605-2496, USA

^gDepartment of Biology, Duke University, Box 90338, Durham, NC 27708, USA

^hDepartment of Plant Biology, University of Minnesota, 1445 Gortner Avenue, St Paul, MN 55108-1095, USA

ⁱDepartment of Biological Sciences, Box 870344/319 Biology, University of Alabama, Tuscaloosa, AL 35487-0344, USA

^jECORC, Agriculture and Agri-Food Canada, CEF, Neatby Building, Ottawa, ON K1A 0C6, Canada

^kDepartment of Botany and Plant Pathology, Oregon State University, 2082 Cordley Hall, Corvallis, OR 97331, USA

^lCentraalbureau voor Schimmelcultures, Uppsalalaan 8, NL-3584 CT Utrecht, The Netherlands

^mUSDA ARS Systematic Botany and Mycology Laboratory, Bldg 011A Rm 319 BARC-WEST, 10300 Baltimore Ave, Beltsville, MD 20705 USA

ⁿABL Herbarium, Gerrit van der Veenstraat 107, NL-3762 XK Soest, The Netherlands

^oBotanical Institute, University of Tübingen, Auf der Morgenstelle 1, D-72076 Tübingen, Germany

^pMax-Planck-Institute for Terrestrial Microbiology, Karl-von-Frisch-Strasse, D-35043 Marburg, Germany

^qDepartment of Plant Pathology, 1453 Fifield Hall, Hull Road, P.O. Box 110680, University of Florida, Gainesville FL 32611-0680, USA

^rInstitute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016, China

^sDepartment of Plant Pathology, Pennsylvania State University, University Park, PA 16802, USA

^tInstitute of Biological Sciences, University of Wales, Aberystwyth SY23 3DA, UK

^uDepartamento de Biología Vegetale II, Facultad de Farmacia, Universidad Complutense de Madrid, Plaza Ramón y Cajal, Ciudad Universitaria, E-28040 Madrid, Spain

* Corresponding author.

E-mail address: dhibbett@clarku.edu (D. S. Hibbett).

0953-7562/\$ – see front matter © 2007 The British Mycological Society. Published by Elsevier Ltd. All rights reserved.

doi:10.1016/j.mycres.2007.03.004

^vDepartment of Biology, University of Oslo, P.O. Box 1066, Blindern, N-0316 Oslo, Norway

^wDepartment of Botany, The Field Museum, 1400 S Lake Shore Dr., Chicago, IL 60605, USA,

^xUSDA-ARS Plant Protection Research Unit, US Plant, Soil and Nutrition Laboratory, Tower Road, Ithaca, NY 14853-2901, USA

^yCentre for Research in Fungal Diversity, Department of Ecology & Biodiversity, University of Hong Kong, Pokfulam Road, Hong Kong SAR, China

^zInstitute of Botany and Ecology, Tartu University, 40 Lai Street, Tartu 51005, Estonia

^{aa}Microbial Properties Research, NCAUR, ARS, USDA, 1815 N. University Street, Peoria, IL 61604-3999, USA

^{ab}Göteborg University, Department of Plant and Environmental Sciences, P. O. Box 461, SE-405 30 Göteborg, Sweden

^{ac}Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66045-7534, USA

^{ad}Department of Biological Sciences, University of Maine, 5722 Deering Hall, Orono, ME 04469-5722, USA

^{ae}Center for Biodiversity, Illinois Natural History Survey, 1816 South Oak Street, Champaign, IL 61820-6970, USA

^{af}Royal Ontario Museum, Centre for Biodiversity and Conservation Biology, 100 Queen's Park, Toronto, ON M5S 2C6, Canada

^{ag}Division of Natural Sciences, Mathematics, and Engineering, Middle Georgia College, 1100 Second Street, SE, Cochran, GA 31014-1599, USA

^{ah}Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, 181 Riia Street Tartu, 51014, Estonia

^{ai}Department of Plant Pathology, Washington State University, P.O. Box 646430, Pullman, WA 99164-6430, USA

^{aj}Chemin des Vignes vieilles, 84120, Mirabeau, France

^{ak}Botanical Institute, University of Oslo, P.O. Box 1066, Blindern, N-0316 Oslo, Norway

^{al}CREM, SABB, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

^{am}Genetics Institute, Department Biology I, Ludwig-Maximilians-University, Maria-Ward-Strasse 1a, D-80638 Munich, Germany

^{an}TechnoSuruga Co. Ltd, Tokyo Office, Ogawamachi Kita Building 4F, 1-8-3, Kanda Ogawamachi, Chiyoda-ku, Tokyo 101-0052, Japan

^{ao}Department of Biology, University of Western Ontario, 213 Biological and Geological Sciences Building, 1151 Richmond Street North, London, ON, N6A 5B7, Canada

^{ap}Department of Systematic Botany, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18 D, SE-752 36, Uppsala, Sweden

^{aq}Botany Department, Brandon University, 270-18th Street, Brandon, MB R7A 6A9, Canada

^{ar}Royal Botanic Garden Edinburgh, 20A Inverleith Row, Edinburgh EH3 5LR, UK

^{as}Faculty of Environmental and Forest Biology, SUNY College, 242 Illick Hall, 1 Forestry Drive, Syracuse, NY 13210, USA

^{at}Department of Biology, Boise State University, 1910 University Drive, S/N Building, Room 210, Boise, ID 83725-1515, USA

^{au}Systematic Mycology and Lichenology Laboratory, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100080, China

^{av}Department of Plant Pathology, Cornell University, Geneva, NY 14456, USA

ARTICLE INFO

Article history:

Received 31 January 2007

Received in revised form

22 February 2007

Accepted 2 March 2007

Published online 13 March 2007

Corresponding Editor:

David L. Hawksworth

Keywords:

AFTOL

Eumycota

Lichens

Molecular phylogenetics

Mycota

Nomenclature

Systematics

ABSTRACT

A comprehensive phylogenetic classification of the kingdom *Fungi* is proposed, with reference to recent molecular phylogenetic analyses, and with input from diverse members of the fungal taxonomic community. The classification includes 195 taxa, down to the level of order, of which 16 are described or validated here: *Dikarya* subkingdom nov.; *Chytridiomycota*, *Neocallimastigomycota* phyla nov.; *Monoblepharidomycetes*, *Neocallimastigomycetes* class. nov.; *Eurotiomycetidae*, *Lecanoromycetidae*, *Mycocaliciomycetidae* subclass. nov.; *Acarosporales*, *Corticiales*, *Baeomycetales*, *Candelariales*, *Gloeophyllales*, *Melanosporales*, *Trechisporales*, *Umbilicariales* orders nov. The clade containing *Ascomycota* and *Basidiomycota* is classified as subkingdom *Dikarya*, reflecting the putative synapomorphy of dikaryotic hyphae. The most dramatic shifts in the classification relative to previous works concern the groups that have traditionally been included in the *Chytridiomycota* and *Zygomycota*. The *Chytridiomycota* is retained in a restricted sense, with *Blastocladiomycota* and *Neocallimastigomycota* representing segregate phyla of flagellated *Fungi*. Taxa traditionally placed in *Zygomycota* are distributed among *Glomeromycota* and several subphyla *incertae sedis*, including *Mucoromycotina*, *Entomophthoromycotina*, *Kickxellomycotina*, and *Zoopagomycotina*. *Microsporidia* are included in the *Fungi*, but no further subdivision of the group is proposed. Several genera of 'basal' *Fungi* of uncertain position are not placed in any higher taxa, including *Basidiobolus*, *Caulochytrium*, *Olpidium*, and *Rozella*.

© 2007 The British Mycological Society. Published by Elsevier Ltd. All rights reserved.

Introduction

The molecular revolution in fungal taxonomy commenced in the early 1990s, with analyses of PCR-amplified ribosomal

RNA genes (White et al. 1990). Today, fungal molecular systematics is a mature discipline in which multi-locus datasets, extensive taxon sampling, and rigorous analytical approaches are standard. To gain an overview of the current state of the

science it is only necessary to survey the recent 'Deep Hypha' issue of *Mycologia* [2007 ('2006'); 98], which contains 21 phylogenetic studies, all of which employ multiple genes to some extent (in some cases, multiple rRNA genes) and that address broad relationships in every major group of *Fungi* (except *Microsporidia*). Another recent milestone is the kingdom-level study of James *et al.* (2006), which used a dataset of six genes (nu-SSU, -LSU, and 5.8S rRNA, *rpb1*, *rpb2* and *tef1*) sampled in nearly 200 species from every major clade of *Fungi* (including *Microsporidia*).

As the broad outlines of fungal phylogeny have come into focus, there have been repeated attempts to summarize the state of knowledge and to restructure higher-level classifications. Two important works that have influenced fungal taxonomy in the 21st century are Ainsworth & Bisby's *Dictionary of the Fungi* (9th edn: Kirk *et al.* 2001), which contains a comprehensive kingdom-wide classification down to the level of genus, and *The Mycota VII* (McLaughlin *et al.* 2001a, 2001b), an edited volume with chapters on all major groups of *Fungi*. These publications represented major advances toward a phylogenetic classification of *Fungi*, but they are already out of date. In the five years since the last edition of the *Dictionary* and the *Mycota VII* appeared, more than 360 articles with the keyword 'phylogen*' were published in *Mycologia* and *Mycological Research* alone, and approximately 80% of the more than 100 000 fungal rRNA gene sequences now in GenBank were deposited (some by molecular ecologists). Recent publications that survey the entire fungal kingdom based on molecular phylogenies include the chapter by Taylor *et al.* (2004) in *Assembling the Tree of Life* (Cracraft & Donoghue 2004), the 'New Higher Level Classification of Eukaryotes' (Adl *et al.* 2005), and the first large collaborative analysis of the *Assembling the Fungal Tree of Life* (AFTOL) project (Lutzoni *et al.* 2004). Taxonomic studies on individual groups of *Fungi* are too numerous to list. Two notable highlights include proposals to recognize the phylum *Glomeromycota* (Schüßler *et al.* 2001) and to include the *Microsporidia* within the *Fungi* (Keeling *et al.* 2000).

On-line fungal taxonomies are also proliferating. One of the most important on-line general classifications of *Fungi* is that of GenBank (www.ncbi.nlm.nih.gov/Taxonomy), which serves a diverse community of researchers, including ecologists and molecular biologists. Another highly visible on-line classification is that of the Tree of Life Web Project (tolweb.org/tree), which is widely used by teachers and students. The classification of *Ascomycota* is being updated regularly via the on-line Myconet series (www.fieldmuseum.org/myconet), and this has been the basis for recent revisions at GenBank, but there is no comparable on-line resource for other major groups of *Fungi*. It is likely that on-line taxonomies will take on even greater prominence in the future, especially as they become integrated with databases of taxonomic names, particularly Index Fungorum (www.indexfungorum.org), MycoBank (www.mycobank.org), and other global biodiversity informatics resources (e.g. Global Biodiversity Information Facility, www.gbif.org).

Although there is broad agreement regarding the composition of the major clades of *Fungi*, there is considerable variation in the names that have been applied to these groups. For example, the clade that is called *Basidiomycetes* in the latest edition of the *Dictionary* is called *Hymenomycetes* at GenBank.

Similarly, the clade that is called *Ascomycetes* in the *Dictionary of the Fungi* is called *Pezizomycotina* in Myconet. Such inconsistencies create confusion, especially for students and non-specialists, and they hamper efforts to develop taxonomic databases.

There is consequently a pressing need for the fungal systematics community to adopt a consensus higher-level classification for the *Fungi* that is based on well-supported monophyletic groups, and which can be recommended for general use. This is an opportune moment to create such a classification. With the new multi-locus analyses, many nodes that were not previously resolved are now supported with confidence. The timing is also good because there are multiple projects in progress that seek to create or update broad classifications of the *Fungi*. In particular, a tenth edition of the *Dictionary* is in preparation, as is a fourth edition of an influential textbook of mycology (Alexopoulos *et al.* 1996). The classifications used by GenBank, the Tree of Life Web Project, and Myconet are being revised continuously. If the classifications employed by these and other major taxonomic resources could be unified, it would promote communication and awareness of fungal phylogeny, and provide a framework for future revisions at all taxonomic levels.

This article presents a higher-level classification for all groups of *Fungi*, with reference to recent molecular phylogenetic studies. The authors represent diverse fungal taxonomy projects, including Ainsworth & Bisby's *Dictionary of the Fungi* (Cannon, Kirk, Stalpers), GenBank (Bischoff), Myconet (Eriksson, Lumbsch, Huhndorf), and Alexopoulos' mycology text (Blackwell, Spatafora). Many of the authors are contributors to the *Fungi* pages in the Tree of Life Web Project. Discussions leading to this classification began in 2004, under the auspices of the AFTOL project and the Deep Hypha Research Coordination Network (Blackwell *et al.* 2007), which were supported by the US National Science Foundation. Throughout the development of this classification, every effort has been made to work in a transparent, consultative manner. The first draft classification was presented at the 2005 Deep Hypha meeting (Tucson, AZ) and subsequently was distributed to a group of 100 fungal systematists for comment. The classification was revised based on comments received and was posted on the AFTOL classification project web site (www.clarku.edu/faculty/dhibbett/AFTOL/AFTOL.htm). Additional modifications were made following the 2006 Deep Hypha meeting (Baton Rouge, LA). For example, the classification of the *Puccinomycotina* was revised to reflect the classification of Bauer *et al.* (2006). The present paper represents a first attempt at a broad-based consensus classification of the *Fungi*. However, the first 20 authors have exercised editorial control and are therefore to be held accountable for errors.

Structure and principles

This classification is restricted to organisms that belong in the monophyletic kingdom *Fungi*, including sexual and asexual forms. It does not consider other organisms formerly included in the kingdom but which are now known not to belong there, even if still studied by mycologists, such as the oomycetes and slime moulds.

The classification adopted here uses a Linnean hierarchy as modified by the *International Code of Botanical Nomenclature (Code)* (McNeill et al. 2006), and uses seven ranks, including: order (suffix: -ales), subclass (-mycetidae), class (-mycetes), subphylum (-mycotina), phylum (-mycota; except *Microsporidia*), subkingdom, and kingdom. The rankings of taxa reflect the preferences and past practices of various authors, as well as the need to keep the nested hierarchies of clades and Linnean categories parallel. Taxa placed at the same rank are not necessarily equivalent in age (except sister taxa), number of species, or degree of morphological divergence.

The classification is limited to taxa down to the level of order. In many orders, especially those representing larger groups, such as *Agaricales*, there is still not enough resolution or taxon sampling to structure a comprehensive family-level classification. The challenge of creating family-level classifications is made even more difficult by the *Code* (McNeill et al. 2006), which requires that names of taxa at the rank of family or lower follow the principle of priority (which does not apply to higher ranks). Ideally, construction of consensus classifications within many of the orders recognized here will involve the coordinated efforts of groups of taxonomic specialists. It is hoped that the present classification will facilitate those endeavors.

The taxa included here are all supported as monophyletic by at least one published phylogenetic analysis (not applicable to monotypic taxa), with the exception of the *Lahmiales* and *Triblidiales* (*Pezizomycotina*) and *Asellariales* (*Kickxellomycotina*), for which molecular data are not available. Support for the monophyly of each group is summarized in three tables, which list selected phylogenetic studies, the type of data that were analysed, the number of OTUs sampled, and BS frequencies and Bayesian PPs. No attempt has been made to cite all of the relevant studies for each group. The analyses chosen for inclusion in the tables are those that have the greatest numbers of loci or taxa, and that provide the strongest support for monophyly of the clades in question. To supplement the information in the tables, brief comments on synonyms, phylogenetic relationships, and composition are provided below for some taxa, along with bibliographic citations for all taxon names. However, it is beyond the scope of this article to discuss each taxon in detail. For additional literature on the phylogeny and taxonomy of individual taxa, readers should consult the studies listed in the tables and below, and the references therein.

The classification is also presented as a set of three tree diagrams. Taxa of uncertain position are listed as *incertae sedis*, and have been placed at the least inclusive level in the hierarchy where they can be assigned with confidence. There are several nodes resolved in the tree figures that are not reflected in the classification. These unnamed clades, for which there is strong to moderate support in recent studies, include the *Dacrymycetes* plus *Agaricomycetes* (*Basidiomycota*) (Matheny et al. 2006, 2007a), *Saccharomycotina* plus *Pezizomycotina* (*Ascomycota*) (James et al. 2006; Spatafora et al. 2007), and the inoperculate euascomycetes (*Ascomycota*) (e.g. Lumbsch et al. 2002). The inoperculate euascomycetes have been recognized as a superclass, the *Leotiomyceta* (Eriksson & Winka 1997; Lumbsch et al. 2002), which is a rank that is not employed here, while the *Dacrymycetes* plus *Agaricomycetes* correspond to the

subclass *Hymenomycetidae* of Swann & Taylor (1995). The absence of these groups from the present classification should not be interpreted as a judgment on their monophyly. Rather, it reflects a desire to keep the classification simple, and to minimize the number of intercalary ranks (as per the directives of Art. 4.3 of the *Code*). Future revisions to this classification will have to consider how to incorporate additional deep nodes, including those that will be resolved with the application of genome-scale datasets (Galagan et al. 2005; Kuramae et al. 2006; Robbertse et al. 2006). One possibility is to employ an unranked category (with or without a uniform suffix) that could be inserted at any level in the taxonomic hierarchy (Hibbett & Donoghue 1998). For example, an unranked classification was adopted in part by Adl et al. (2005).

Overview of the classification

The classification accepts one kingdom, one subkingdom, seven phyla, ten subphyla, 35 classes, 12 subclasses, and 129 orders. Taxa that are described or validated here include *Chytridiomycota*, *Monoblepharidomycetes*, *Neocallimastigomycota*, *Neocallimastigomycetes*, *Dikarya*, *Acarosporales*, *Baeomycetales*, *Candelariales*, *Umbilicariales*, *Lecanoromycetidae*, *Eurotiomycetidae*, *Mycocaliciomycetidae*, *Melanosporales*, *Corticiales*, *Gloeophyllales*, and *Trechisporales*. Thus, about 90% of the 195 taxon names employed in the present classification have been validly published previously. The clade containing the *Ascomycota* and *Basidiomycota* is classified as the subkingdom *Dikarya* (as used in James et al. 2006), reflecting the putative synapomorphy of dikaryotic hyphae (Tehler 1988). All of the other new names are based on automatically typified teleomorphic names. The classification of *Ascomycota* largely parallels that of the Myconet classification, including recent changes that will be adopted in the forthcoming 2007 'Outline of the *Ascomycota*'. In *Basidiomycota*, the clades formerly called *Basidiomycetes*, *Urediniomycetes*, and *Ustilaginomycetes* in the last edition of Ainsworth & Bisby's *Dictionary of the Fungi* are called the *Agaricomycotina*, *Pucciniomycotina*, and *Ustilaginomycotina*, respectively, as in Bauer et al. (2006). This is done to minimize confusion between taxon names and informal terms (*basidiomycetes* is a commonly used informal term for all *Basidiomycota*) and to refer to the included genera *Agaricus* (including the cultivated button mushroom) and *Puccinia* (which includes barberry-wheat rust). Another significant change in the *Basidiomycota* classification is the inclusion of the *Wallemiomycetes* and *Entorrhizomycetes* as classes *incertae sedis* within the phylum, reflecting ambiguity about their higher-level placements (Matheny et al. 2007b).

The most dramatic changes in the classification concern the 'basal fungal lineages', which include the taxa that have traditionally been placed in the *Zygomycota* and *Chytridiomycota*. These groups have long been recognized to be polyphyletic, based on analyses of rRNA, *tef1*, and *rpb1* (James et al. 2000; Nagahama et al. 1995; Tanabe et al. 2004, 2005). The recent multilocus analyses of James et al. (2006) and others now provide the sampling, resolution, and support necessary to structure new classifications of these early-diverging groups, although significant questions remain. The *Chytridiomycota* is retained in a highly restricted sense, including

Chytridiomycetes and Monoblepharidomycetes. The Blastocladias, a traditional member of the Chytridiomycota, is here treated as a phylum, the Blastocladiomycota, as in James *et al.* (2007). The Neocallimastigales, whose distinctiveness from other chytrids has long been recognized, is also elevated to phylum, based on both morphology and molecular phylogeny. The genera *Caulochytrium*, *Olpidium*, and *Rozella*, which have traditionally been placed in the Chytridiomycota, and *Basidiobolus*, which has been classified in the Zygomycota (Entomophthorales), are not included in any higher taxa in this classification, pending more definitive resolutions of their placements.

The phylum Zygomycota is not accepted in this classification, pending resolution of relationships among the clades that have traditionally been placed in the Zygomycota (see discussion under Mucoromycotina). The traditional Zygomycota are here distributed among the phylum Glomeromycota and four subphyla *incertae sedis*, including Mucoromycotina, Kickxellomycotina, Zoopagomycotina and Entomophthoromycotina. A clade containing the Glomeromycota and the Dikarya was resolved previously based on ribosomal RNA genes and was classified as the Symbiomyota (Tehler *et al.* 2003). That taxon is not included here, because there was not strong support for the clade in the analyses of James *et al.* (2006) or Liu *et al.* (2006). If the Symbiomyota is added to this classification, it will need to be assigned a rank between kingdom and subkingdom, or perhaps be classified as an unranked taxon.

Microsporidia, unicellular parasites of animals and protists with highly reduced mitochondria (Germot *et al.* 1997; Hirt *et al.* 1997; Peyretilade *et al.* 1998), are included here as a phylum of the Fungi, based on analyses by Keeling *et al.* (2000), Gill & Fast (2006), James *et al.* (2006), and Liu *et al.* (2006). The latter study concluded that Microsporidia are the sister group of the rest of the Fungi and should not be classified as true Fungi, but that topology does not conflict with the delimitation of the monophyletic Fungi as proposed here. The analysis of James *et al.* (2006) suggested that *Rozella*, which was not sampled by Liu *et al.* (2006), is the sister group of the Microsporidia. No subdivision of the Microsporidia is proposed, owing to a lack of well-sampled multilocus analyses of this group (but see Vossbrinck & Debrunner-Vossbrinck 2005, for an analysis using SSU rRNA genes).

Phylogenetic classification of Fungi

Many of the citations and authorities in the list below were obtained from the Index Fungorum databases (www.indexfungorum.org). A brief list of exemplar genera, including the type for automatically typified names, is given for each order (for small orders, all included genera are listed). A number of the genera listed are used in a modern, restricted sense, and readers are urged to consult the primary literature cited below and in the tables for information about current generic concepts. Comprehensive lists of genera and families included in each order will be forthcoming in the *Dictionary of the Fungi* (10th edn; listing on-line at www.indexfungorum.org) and in the next revision of Myconet (for Ascomycota). Further information on the names of fungi (not only kingdom

Fungi) above the rank of order and their places of publication may be found in the preliminary catalogue compiled by David (2002).

In accordance with the practice in recent editions of the Code, all scientific names regardless of rank are placed in italic type here except in the first line of the treatment of each accepted taxon where they are given in bold Roman type to make them stand out. When these names are used by other mycologists in their own publications, we wish to encourage the practice of the use of italics as recommended in the Preface to the current Code (McNeill *et al.* 2006).

Kingdom: **Fungi** R. T. Moore, *Bot. Mar.* 23: 371 (1980).

Synonym: Fungi T. L. Jahn & F. F. Jahn, *How to Know the Protozoa*: 7 (1949), *nomen invalidum*. (Table 1, Fig 1)

The concept of the Fungi as one of six kingdoms of life was introduced by Jahn & Jahn (1949), and a five kingdom system was advanced by Whittaker (1959), but neither of these works included a Latin diagnosis and the name was therefore invalid under the Code until the required Latin was provided by Moore (1980). Although Moore did not make a specific reference to Jahn & Jahn's book, he was well aware that the name was in widespread use in the rank of kingdom. Under the current Code, Jahn & Jahn are not to be included in the author citation. However, a proposal to change this provision in the Code will be made at the next International Botanical Congress (D. L. Hawksworth, unpubl.). If it is approved, the correct citation would be Fungi T. L. Jahn & F. F. Jahn ex R. T. Moore (this rule change would also affect the citations of Ascomycota and Basidiomycota).

Phylum: **Chytridiomycota** M. J. Powell, **phylum nov.**

Mycobank no.: MB 501278

Synonyms: *Archemycota* Caval.-Sm., *Biol. Rev.* 73: 246 (1998), *pro parte*.

Thallus monocentricus vel polycentricus vel filamentosus; propagatio asexualis zoosporis, flagello retrorsum inserto, kinetosomate et centriolo supervacaneo praeditis, 9 munimentis flagelli, et complexu "microbody-corpore lipideo" descriptis; propagatio sexualis meiosi post copulationem perfecta; apparatus Golgi e cisternis superimpositis constans; tegumentum nuclei mitosi procedente circum polos fenestratum.

Typus: *Chytridium* A. Braun 1851.

Thallus monocentric, polycentric, or filamentous; asexual reproduction by zoospores with a single posteriorly-directed flagellum, both a kinetosome and non-functional centriole, nine flagellar props, and a microbody-lipid globule complex; sexual reproduction with zygotic meiosis where known; Golgi apparatus with stacked cisternae; nuclear envelope fenestrated at poles during mitosis.

Used as a phylum name without Latin diagnosis or description among others by von Arx (1967) and Margulis *et al.* (1990). Equivalent to euchytrids of James *et al.* 2006, the 'core chytrid clade' of James *et al.* (2007), or the 'core chytrid clade' plus the *Monoblepharidales* of James *et al.* (2000). Earlier usages are not indicated in the author citation of the name, because the circumscription adopted here differs significantly from that of those authors.

Table 1 – Support for major groups of Fungi in selected phylogenetic studies: basal fungi and Dikarya

Rank	Taxon	Reference	Data ^a	OTUs ^b	Support ^c
Kingdom	FUNGI	Keeling (2003)	α -tub, β -tub	38	MLBS = 98 NJBS = 94
		Baldauf et al. (2000)	act, α -tub, β -tub, tef1	12	MLBS = 85 MPBS = 95
Phylum	CHYTRIDIOMYCOTA	James et al. (2007)	LSU, SSU, 5.8S	84	BPP \geq 0.95
		Seif et al. (2005)	mt-genome	5	BPP = 1
Class	Chytridiomycetes	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	8	MLBS = 100 BPP \geq 0.95
		James et al. (2007)	LSU, SSU, 5.8S	75	MLBS \geq 70 BPP \geq 0.95
		Keeling (2003)	α -tub, β -tub	5	MLBS = 90 NJBS = 95
Order	Chytridiales	James et al. (unpublished)	LSU, SSU, 5.8S, rpb1, rpb2, tef1, atp6	9	MLBS = 98
Order	Rhizophydiales	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	2	BPP \geq 0.95 MLBS \geq 70
		Letcher et al. (2006)	LSU, 5.8S	96	MPBS = 100 BPP = 1
Order	Spizellomycetales	James et al. (2007)	LSU, SSU, 5.8S	9	MPBS = 100
Class/Order	Monoblepharidomycetes, Monoblepharidales	James et al. (2007)	LSU, SSU, 5.8S	9	BPP \geq 0.95 MLBS \geq 70 MPBS \geq 70
Phylum/Class/Order	NEOCALLIMASTIGOMYCOTA, Neocallimastigomycetes, Neocallimastigales	Bullerwell et al. (2003) James et al. (2007)	cox 1,2,3; cob, atp6,9; nad 1,2,3,4, 4L,6 LSU, SSU, 5.8S	4 6	MLBS = 100 BPP \geq 0.95 MLBS \geq 70 MPBS \geq 70
Phylum/Class/Order	BLASTOCLADIOMYCOTA, Blastocladiomycetes, Blastocladales	James et al. (2007)	LSU, SSU, 5.8S	10	BPP \geq 0.95
		Liu et al. (2006)	rpb1, rpb2	3	BPP = 1 MPBS = 100
Phylum	MICROSPORIDIA	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	2	BPP \geq 0.95 MLBS \geq 70
		Keeling (2003)	α -tub, β -tub	6	MLBS = 100 NJBS = 97
Phylum/Class	GLOMEROMYCOTA, Glomeromycetes	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	5	BPP \geq 0.95 MLBS \geq 70
Order	Archaeosporales	Schüßler et al. (2001)	SSU	72	NJBS \geq 90
Order	Diversisporales	Schüßler et al. (2001)	SSU	5	NJBS \geq 95
Order	Glomerales	Schüßler et al. (2001)	SSU	32	NJBS \geq 95
Order	Paraglomerales	Schüßler et al. (2001)	SSU	32	NJBS \geq 95
	Subphyla incertae sedis (not placed in any phylum)			3	NJBS \geq 95
Subphylum	Mucoromycotina	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	11	BPP = 1
		Tanabe et al. (2004)	rpb1	4	NJBS = 82
Order	Mucorales	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	3	BPP \geq 0.95 MLBS \geq 70
		Tanabe et al. (2004)	rpb1	3	NJBS = 100
		Keeling (2003)	α -tub, β -tub	4	MLBS = 96 NJBS = 98
		White et al. (2007)	LSU, SSU, 5.8S	28	BPP = 1 MPBS \geq 70
Order	Endogonales	White et al. (2007)	LSU, SSU, 5.8S	2	BPP = 1 MPBS \geq 70
Order	Mortierellales	White et al. (2007)	LSU, SSU, 5.8S	6	BPP = 1 MPBS \geq 70
Subphylum/Order	Entomophthoromycotina, Entomophthorales	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	2	BPP \geq 0.95 MLBS \geq 70
Subphylum/Order	Zoopagomycotina, Zoopagales	James et al. (2006)	LSU, SSU, 5.8S, rpb1, rpb2, tef1	2	BPP \geq 0.95 MLBS \geq 70
		Tanabe et al. (2004)	rpb1	3	NJBS = 86
Subphylum	Kickxellomycotina	Tanabe et al. (2004)	rpb1	6	NJBS = 84
Order	Kickxellales	O'Donnell et al. (1998)	SSU	7	MPBS = 100
Order	Dimargaritales	Tanabe et al. (2000)	SSU	3	NJBS = 100

Table 1 (continued)

Rank	Taxon	Reference	Data ^a	OTUs ^b	Support ^c
Order	<i>Harpellales</i>	Tanabe <i>et al.</i> (2004)	<i>rpb1</i>	3	NJBS = 98
		O'Donnell <i>et al.</i> (1998)	SSU	4	MPBS = 100
Order	<i>Asellariales</i>	—	—	—	—
Subkingdom	DIKARYA	James <i>et al.</i> (2006)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	161	BPP = 1 MLBS = 71
		Steenkamp <i>et al.</i> (2006)	<i>act</i> , α - <i>tub</i> , β - <i>tub</i> , <i>tef1</i>	10	BPP = 1 MLBS = 84 MPBS = 82 NJBS = 96
		Seif <i>et al.</i> (2005)	mt-genome	10	BPP = 1 MLBS = 100
		Liu <i>et al.</i> (2006)	<i>rpb1</i> , <i>rpb2</i>	27	BPP = 1 MPBS = 100

Taxa with only one subsidiary taxon included (i.e. redundant taxa) are listed on a single line, with rank abbreviations divided by a slash (e.g. the class *Agaricostilbomycetes*, which contains a single order, *Agaricostilbales*, is indicated as Class/Order).

a LSU, SSU, and 5.8S refer to nuclear rRNA genes, whereas mt-LSU and mt-SSU refer to mitochondrial rRNA genes, other genes follow standard abbreviations. Some datasets contain missing sequences.

b Indicates the number of OTUs in the specified clade, not the total number of OTUs in the dataset.

c BS, bootstrap %, jk, jackknife %, WP = weighted parsimony, RML = RaxML, PML = PhyML, ME = minimum evolution, BPP, Bayesian posterior probability, NA, not applicable because the group is monotypic, or only a single species was sampled in the reference study.

Class: **Chytridiomycetes** Caval.-Sm., *Biol. Rev.* 73: 246 (1998).
 Synonym: *Archimycetes* A. Fisch. (Fischer 1892) *pro parte*
 (included *Olpidiopsis*, *Hypochoytrium*).
 Type: *Chytridium* A. Braun 1851.

Reproducing asexually by zoospores bearing a single posteriorly-directed flagellum; zoospores containing a kinetosome




Fig 1 – Phylogeny and classification of Fungi. Basal Fungi and Dikarya. Branch lengths are not proportional to genetic distances. See Table 1 for support values for clades.

and a non-flagellated centriole; thallus monocentric or rhizomycelial polycentric; sexual reproduction not oogamous.

Cavalier-Smith (1998) provided a brief, four-word, Latin description that was not diagnostic for phyla of uniflagellate fungi, and has been revised above. The name *Chytridiomycetes* was also used by Serbinow (1907), Cejp (1957), Sparrow (1958), and Alexopoulos *et al.* (1996). For further discussion of the nomenclatural history of the name, see David (2002).

Order: **Chytridiales** Cohn, *Jber. schles. Ges. vaterl. Kultur* 57: 279 (1879).

Emend. Schröter (as 'Chytridineae') in Engler & Prantl, *Nat. Pflanzenfam.* 1: 64 (1892). Emend. Barr, *Can. J. Bot.* 58: 2384 (1980). Emend. Letcher & Powell, *Mycol. Res.* 110: 907 (2006).

Type: *Chytridium* A. Braun 1851.

Thallus monocentric or polycentric rhizomycelial; zoospores typically with flagellar base containing an electron-opaque plug, microtubules extending from one side of the kinetosome in a parallel array, ribosomes aggregated near the nucleus, kinetosome parallel to non-flagellated centriole and connected to it by fibrous material, nucleus not associated with kinetosome, fenestrated cisterna (rumposome) adjacent to lipid globule.

Exemplar genera: *Chytridium* A. Braun 1851, *Chytriomycetes* Karling 1945, *Nowakowskiella* J. Schröt. 1893.

An emended description is presented above to conform to the circumscription adopted here. Monophyly of this group, as currently delimited, is not certain; *Polychytrium* Ajello 1942 and its allies and *Chytriomycetes angularis* Longcore 1992 and its allies may eventually be segregated from *Chytridiales* s. str.

Order: **Rhizophydiales** Letcher, in Letcher *et al.*, *Mycol. Res.* 110: 908 (2006).

Exemplar genera: *Rhizophyidium* Schenk 1858, *Kappamyces* Letcher & M.J. Powell 2005, *Terramyces* Letcher 2006,

Boothomyces Letcher 2006; *Batrachochytrium* Longcore, Pessier & D.K. Nichols 1999 is on a long branch in this clade with no near relatives.

Order: **Spizellomycetales** D. J. S. Barr, *Can. J. Bot.* **58**: 2384 (1980).

Exemplar genera: *Spizellomyces* D. J. S. Barr 1980, *Powellomyces* Longcore, D. J. S. Barr & Désauln. 1995, *Kochiomyces* D. J. S. Barr 1980.

This classification does not include *Caulochytrium*, *Olpidium*, *Rozella*, or the *Rhizophlyctis rosea* clade, which are considered *incertae sedis*.

Class: **Monoblepharidomycetes** J. H. Schaffner, *Ohio Nat.* **9**: 449 (1909), as '*Monoblepharideae*'.

Type: *Monoblepharis* Cornu 1871.

Thallus filamentous, either extensive or a simple unbranched thallus, often with a basal holdfast; asexual reproduction by zoospores or autospores; zoospores containing a kinetosome parallel to a non-flagellated centriole, a striated disk partially extending around the kinetosome, microtubules radiating anteriorly from the striated disk, a ribosomal aggregation, and rumposome (fenestrated cisterna) adjacent to a microbody; sexual reproduction oogamous by means of posteriorly uniflagellate antherozoids borne in antheridia and nonflagellate female gametes borne in oogonia.

Schaffner (1909) used the name '*Monoblepharideae*' as a class but with the ending of a suborder; this must be changed without change of authorship or date of publication (*Code*, Art. 16.3).

Order: **Monoblepharidales** J. Schröt., in Engler & Prantl, *Nat. Pflanzenfam.* **1**: 106 (1893), as '*Monoblepharidinae*'.

Emend. Sparrow, *Aquatic Phycomycetes*: 458 (1943).

Emended description as for *Monoblepharidomycetes*.

Exemplar genera: *Monoblepharis* Cornu 1871, *Harpochytrium* Lagerh. 1890, *Oedogoniomyces* Tak. Kobay. & M. Ôkubo 1954.

Phylum: **Neocallimastigomycota** M. J. Powell, **phylum nov.**

Mycobank no.: MB 501279

Thallus monocentricus vel polycentricus; fungi anaerobici, intra tractum digestivum animalium herbivororum vel fortasse in substratis anaerobicis terrestribus vel limnicis; mitochondriis carentes sed hydrogenosomatibus praediti; zoosporae retrorsum uni- vel multiflagellatae, kinetosoma praesens sed centriolum supervacaneum absens; complexus kinetosomati affixus e radio marginali et annulo circumflagellari compositus; microtubuli e radio entendentis circum nucleum radiantes et flabellum posterius formantes; munimenta flagelli absentia; tegumentum nuclei mitosi procedente integrum remanens.

Typus: *Neocallimastix* Vavra & Joyon ex I. B. Heath 1983.

Thallus monocentric or polycentric; anaerobic, found in digestive system of larger herbivorous mammals and possibly in other terrestrial and aquatic anaerobic environments; lacks mitochondria but contains hydrogenosomes of mitochondrial origin; zoospores posteriorly unflagellate or polyflagellate, kinetosome present but non-functional centriole absent, kinetosome-associated complex composed of a skirt, strut, spur and circumflagellar ring, microtubules extend from spur and radiate around nucleus, forming a posterior fan, flagellar

props absent; nuclear envelope remains intact throughout mitosis.

Class: **Neocallimastigomycetes** M. J. Powell, **class. nov.**

Mycobank no.: MB 501280

Diagnosis latina ut in *Neocallimastigomycota* (vide supra).

Typus: *Neocallimastix* Vavra & Joyon ex I.B. Heath 1983.

Order: **Neocallimastigales** J. L. Li, I. B. Heath & L. Packer, *Can. J. Bot.* **71**: 403 (1993).

Exemplar genera: *Neocallimastix* Vavra & Joyon ex I.B. Heath 1983, *Caecomycetes* J.J. Gold 1988, *Orpinomyces* D.J.S. Barr, H. Kudo, Jakober & K.J. Cheng 1989.

Phylum: **Blastocladiomycota** T. Y. James, *Mycologia* **98**: 867 (2007) ['2006'].

Synonym: *Allomycota* Caval.-Sm., *BioSystems* **14**: 465 (1981).

This phylum was proposed to reflect phylogenetic information from a number of molecular studies (James et al. 2007; Liu et al. 2006).

Class: **Blastocladiomycetes** T. Y. James, *Mycologia* **98**: 868 (2007) ['2006'].

Synonym: *Allomycetes* Caval.-Sm., *Biol. Rev.* **73**: 246 (1998), based on *Allomyces* E. J. Butler 1911.

Cavalier-Smith provided a brief, five-word Latin description for *Allomycetes* that is not diagnostic from other uniflagellate fungi. The name *Allomycetes* was not taken up, because it is appropriate to have a class name based on the same genus as an included ordinal name, and because Cavalier-Smith's 'diagnosis' was vague.

Order: **Blastocladiiales** 1910, H. E. Petersen, *Bot. Tidsskr.* **29**: 357 (1909) ('*Blastocladiinae*').

Exemplar genera: *Allomyces* E. J. Butler 1911, *Blastocladia* Reinsch 1877, *Coelomomyces* Keilin. 1921.

Phylum: **Microsporidia** Balbiani, *C. R. Acad. Sci. Paris* **95**: 1168 (1882).

The nomenclatural status of *Microsporidia* is ambiguous. It has been treated as a phylum under the zoological *Code* (International Commission on Zoological Nomenclature 1999), but there is disagreement about the correct author citation (Larsson 2000; Sprague & Becnel 1998), and it is uncertain if the name would be valid under the botanical *Code*. This uncertainty arises as *Microsporidium* Balbiani 1884 appears to be a later synonym of *Nosema* Naegeli 1857. The present work follows the recommendation of Sprague & Becnel (1998) in attributing *Microsporidia* to Balbiani (1882), but this must be regarded as provisional. Before the status of the *Microsporidia* can be resolved, it will be necessary to decide whether the nomenclature of the group as a whole should be governed by the zoological or the botanical *Code* although the latter now allows names of fungi described under the zoological *Code* to be accepted. The final decision will require input from the community of scientists who study *Microsporidia*.

No subdivision of the group is proposed here, owing to the lack of well-sampled multi-gene phylogenies within the group. However, Vossbrinck & Debrunner-Vossbrinck (2005) proposed a class-level classification of microsporidia, based on small-subunit rRNA gene sequences.

Phylum: **Glomeromycota** C. Walker & A. Schuessler, in Schüßler et al., *Mycol. Res.* **105**: 1416 (2001).

Class: **Glomeromycetes** Caval.-Sm., *Biol. Rev.* **73**: 246 (1998), as 'Glomomycetes'.

Synonym: *Geomyces* Caval.-Sm., *Biol. Rev.* **73**: 247 (1998).

Order: **Archaeosporales** C. Walker & A. Schuessler, in Schüßler et al., *Mycol. Res.* **105**: 1418 (2001).

Synonym: *Geosiphonales* Caval.-Sm., *Biol. Rev.* **73**: 247 (1998).

Exemplar genera: *Archaeospora* J.B. Morton & D. Redecker 2001, *Geosiphon* F. Wettst. 1915.

Order: **Diversisporales** C. Walker & A. Schuessler, *Mycol. Res.* **108**: 981 (2004).

Exemplar genera: *Acaulospora* Gerd. & Trappe 1974, *Diversispora* C. Walker & A. Schüßler 2004, *Gigaspora* Gerd. & Trappe 1974, *Pacispora* Oehl & Sieverd. 2004.

Order: **Glomerales** J. B. Morton & Benny, *Mycotaxon* **37**: 473 (1990), as 'Glomales'.

Exemplar genus: *Glomus* Tul. & C. Tul. 1845.

Order: **Paraglomerales** C. Walker & A. Schuessler, in Schüßler et al., *Mycol. Res.* **105**: 1418 (2001).

Exemplar genus: *Paraglomus* J. B. Morton & D. Redecker 2001.

Subphyla incertae sedis (not assigned to any phylum):

Subphylum: **Mucoromycotina** Benny, **subphylum nov.**

Mycobank no.: MB 501281

Fungi saprotrophici vel raro mycoparasiti facultativi, gallas facientes, haustoriis carentes, raro ectomycorrhizam facientes. Mycelium ramosum, juvene coenocyticum, maturum aliquando septis microporosis divisum. Reproductio asexualis sporangiis vel sporangiolis vel merosporangiis, raro chlamydozporis vel arthrosporis vel blastozporis effecta. Reproductio sexualis zygosporis plus minusve globosis e suspensoribus oppositis vel appositis formatis effecta.

Typus: *Mucor* Fresen. 1850.

Fungi saprobes, or rarely gall-forming, nonhaustorial, facultative mycoparasites, or forming ectomycorrhiza. Mycelium branched, coenocytic when young, sometimes producing septa that contain micropores at maturity. Asexual reproduction by sporangia, sporangiola, or merosporangia, or rarely by chlamydozspores, arthrospores, or blastospores. Sexual reproduction by more or less globose zygosporis formed on opposed or apposed suspensors.

This group includes the *Mucorales*, which is the core group of the traditional *Zygomycota*. Monophyly of the traditional *Zygomycota* (including *Mucorales*, *Glomerales*, *Entomophthorales* and *Harpellales*) was suggested by a recent study by Liu et al. (2006) using *rpb1* and *rpb2*, but that finding conflicts with results of

analyses that included additional loci and taxa, which suggested that the traditional *Zygomycota* is polyphyletic (James et al. 2006).

The name *Zygomycota* was first published without a Latin diagnosis by Moreau (1954) and is therefore invalid. At present, this classification does not include *Zygomycota*. When relationships among basal fungal lineages are more clearly resolved, it may be appropriate to resurrect and validate *Zygomycota*, to include *Mucoromycotina* and perhaps other clades.

Order: **Mucorales** Fr., *Syst. Mycol.* **3** (2): 296 (1832).

Exemplar genera: *Mucor* Fresen. 1850 (pro parte), *Parasitella* Bainier 1903, *Phycomyces* Kunze 1823, *Pilobolus* Tode 1784, *Rhizopus* Ehrenb. 1821.

Order: **Endogonales** Moreau ex R. K. Benj., in Kendrick (ed.), *Whole Fungus* **2**: 599 (1979).

Emend.: Morton & Benny, *Mycotaxon* **37**: 473 (1990).

Synonym: *Endogonales* Moreau, *Encycl. Mycol.* **23**: 1231 (1954), *nomen invalidum*.

Exemplar genera: *Endogone* Link 1809, *Peridiospora* C. G. Wu & S. J. Lin 1997, *Sclerogone* Warcup 1990, *Youngiomyces* Y. J. Yao 1995.

Order: **Mortierellales** Caval.-Sm., *Biol. Rev.* **73**: 246 (1998).

Exemplar genera: *Mortierella* Coem. 1863, *Dissophora* Thaxt. 1914, *Modicella* Kanouse 1936.

Subphylum: **Entomophthoromycotina** Humber, **subphylum nov.**

Mycobank no.: MB 501282

Fungi pathogenici obligate animalibus (praecipue invertebratis) vel plantis cryptogamicis vel saprotrophicis, interdum in animalibus vertebratis parasitici. Status somaticus mycelium coenocyticum vel septatum, pariete circumdatum vel protoplasticum, in hospite culturisve saepe corpora hyphalia multinucleata formans; forma protoplastica hyphoidea vel amoeboida forma variabilis; cystidia et rhizoidea in aliquot speciebus athropodocolis formata. Characteres nuclei, sicut magnitudo, nucleoli magnitudo et locus, praesentia aut absentia heterochromatini intermitotici, familiis distinguendis iuvant. Conidiophora simplicia ramosave. Sporae primariae conidia vera, uninucleatae vel plurinucleatae vel multinucleatae, variis modis vi propulsae vel passive liberatae, conidia secundaria persaepe formata. Sporae perdurantes crassituncatae, bistratosae velut zygosporae post conjugationem velut azygosporae singulae formatae.

Typus: *Entomophthora* Fresen. 1856.

Obligate pathogens of animals (primarily arthropods), cryptogamic plants, or saprobes; occasionally facultative parasites of vertebrates. Somatic state consisting of a well-defined mycelium, coenocytic or septate, walled or protoplastic, which may fragment to form multinucleate hyphal bodies; protoplasts either hyphoid or amoeboid and changeable in shape; cystidia or rhizoids formed by some taxa. Such nuclear characters as overall size, location and comparative size of nucleoli, presence or absence of granular heterochromatin in chemically unfixed interphasic nuclei, and mitotic patterns are important at the family level. Conidiophores branched or unbranched. Primary spores true conidia, uni-, pluri-, or multinucleate, forcibly

discharged by diverse possible means or passively dispersed; secondary conidia often produced. Resting spores with thick bi-layered walls form as zygospores after conjugations of undifferentiated gametangia from different or the same hyphal bodies or hypha or as azygospores arising without prior gametangial conjugations.

Order: **Entomophthorales** G. Winter, *Rabenh. Krypt.-Fl.* 1: 74 (1880).

Exemplar genera: *Entomophthora* Fresen. 1856, *Ballocephala* Drechsler 1951, *Conidiobolus* Bref. 1884, *Entomophaga* Batko 1964, *Neozygites* Witlaczil 1885.

Subphylum: **Zoopagomycotina** Benny, **subphylum nov.**

Mycobank no.: MB 501283

Fungi endo- vel ectoparasitici microanimalium vel fungorum. Corpus vegetativum ex thallo simplici ramoso vel nonramoso vel mycelio nonseptato plus minusve extense ramoso constans. Ectoparasitae haustoria intra hospitem formantes. Reproductio asexualis arthrosporis, chlamydosporis vel sporangiolis uni- vel multisporis perfecta; sporangiosporae sporangiolorum multisporum in catenensis (merosporangiis) simplicibus vel ramosis dispositae. Reproductio sexualis zygosporis paene globosis perficitur; hyphae sexuales hyphis vegetativis similes vel plus minusve ampliatae.

Typus: *Zoopage* Drechsler 1935.

Endo- or ectoparasites of microanimals and fungi. Vegetative body consisting of a simple, branched or unbranched thallus or more of less extensively branched mycelium. Ectoparasites forming haustoria inside the host. Asexual reproduction by arthrospores, chlamydospores or uni- or multispored sporangia; sporangiospores of multispored sporangia formed in simple or branched chains (merosporangia). Sexual reproduction by nearly globose zygospores; sexual hyphae similar to the vegetative hyphae or more or less enlarged.

The description of this group is based mostly on the validating description for the *Zoopagales* by Benjamin (1979), except that arthrospores have been added, based on Barron's (1975) report of arthrospores in *Helicocephalum* Thaxt. 1891.

Order: **Zoopagales** Bessey ex R.K. Benj., in Kendrick (ed.), *Whole Fungus* 2: 590 (1979).

Synonym: *Zoopagales* Bessey, *Morph. Tax. Fungi* : 177 (1950), *nomen invalidum*.

Exemplar genera: *Cochlonema* Drechsler 1935, *Rhopalomyces* Corda 1839, *Piptocephalis* de Bary 1865, *Sigmoideomyces* Thaxt. 1891, *Syncephalis* Tiegh. & G. Le Monn. 1873, *Zoopage* Drechsler 1935.

Subphylum: **Kickxellomycotina** Benny, **subphylum nov.**

Mycobank no.: MB 501284

Fungi saprotrophici vel mycoparasitici vel obligate symbiotici. Thallus in nonnullis generibus e tenaculo fungos alios parasitans et haustoriis penetrans; mycelium septatum, ramosum vel simplex; septa in medio excavata et obturata. Reproductio asexualis merosporangiis uni- vel bisporis vel trichosporis vel arthrosporis effecta. Reproductio sexualis zygosporis globosis, biconicis vel allantoidis circinatis effecta.

Typus: *Kickxella* Coem. 1862.

Fungi saprobes, mycoparasites, or obligate symbionts. Thallus arising from a holdfast on other fungi as a haustorial parasite, or branched, septate, subaerial hyphae. Mycelium branched or unbranched, regularly septate. Septa with median, disciform cavities containing plugs. Asexual production by 1- or 2-spored merosporangia, trichospores, or arthrospores. Sexual reproduction by zygospores that are globose, biconical, or allantoid and coiled.

Order: **Kickxellales** Kreisel ex R. K. Benj., in Kendrick (ed.), *Whole Fungus* 2: 610 (1979).

Synonym: *Kickxellales* Kreisel, *Grundz. nat. Syst. Pilze*: 65 (1969), *nomen invalidum*.

Exemplar genera: *Kickxella* Coem. 1862, *Coemansia* Tiegh. & G. Le Monn. 1873, *Linderina* Raper & Fennell 1952, *Spirodactylon* R. K. Benj. 1959.

Order: **Dimargaritales** R. K. Benj., in Kendrick (ed.), *Whole Fungus* 2: 607 (1979).

Exemplar genera: *Dimargaris* Tiegh. 1875, *Dispira* Tiegh. 1875, *Tieghemiomyces* R. K. Benj. 1959.

Order: **Harpellales** Lichtw. & Manier, *Mycotaxon* 7: 441 (1978).

The taxa in this order have been referred to as 'Trichomycetes'. However, *Trichomycetes* is no longer a useful phylogenetic taxon because it describes a polyphyletic group. The use of the term should be restricted to ecological rather than phylogenetic groupings, and not capitalized or italicized, i.e. as 'trichomycetes'.

Exemplar genera: *Harpella* L. Léger & Duboscq 1929, *Furculomyces* Lichtw. & M. C. Williams 1992, *Legeriomyces* Pouzar 1972, *Smittium* R. Poiss. 1937.

Order: **Asellariales** Manier ex Manier & Lichtw., *Mycotaxon* 7: 442 (1978).

Exemplar genera: *Asellaria* R. Poiss. 1932, *Orchesellaria* Manier ex Manier & Lichtw. 1968.

Asellariales are retained in the *Fungi* here due to their ultrastructural characteristics (Benny & White 2001; Manier 1973; Moss 1975; Saikawa et al. 1997). Unpublished *rpb1* and *rpb2* data also support their placement in the *Kickxellomycotina* (T. Y. James & M. M. White, unpubl.).

Subkingdom: **Dikarya** Hibbett, T. Y. James & Vilgalys, **subregnum nov.**

Mycobank no.: MB 501285

Synonyms: *Neomycota* Caval.-Sm., *Rev. Biol.* 73: 209 (1998).

Carpomycetaceae Bessey, *Univ. Studies, Univ. Nebr.* 7: 294 (1907).

Fungi unicellulares vel filamentosi, flagellis carentes, saepe stadium dikaryoticum includentes. *Ascomycota* et *Basidiomycota* complectens.

Unicellular or filamentous *Fungi*, lacking flagella, often with a dikaryotic state. The least-inclusive clade that contains *Ascomycota* and *Basidiomycota*.

The name alludes to the putative synapomorphy of dikaryotic hyphae (Tehler 1988) and was applied by James *et al.* (2006) without formal description. Kendrick (1985) and Tehler *et al.* (2003) referred to this group as the Dikaryomycota, but the termination ‘-mycota’ denotes the rank of phylum under the Code. Cavalier-Smith (1998) referred to this group as Neomycota. Dikarya is used here, because it is more descriptive and is consistent with recent use (James *et al.* 2006; Tehler *et al.* 2003; Kendrick 1985).

Phylum: **Ascomycota** Caval.-Sm., *Biol. Rev.* 73: 247 (1998), as ‘Ascomycota Berk. 1857. stat. nov.’

Synonyms: *Ascomycetes* Berk., *Intr. Crypt. Bot.*: 270 (1857), rank uncertain; Whittaker (1959: 220).

Ascomycota Bold, *Morph. Pl.*: 7, 180 (1958), *nomen invalidum*; Hawksworth *et al.* (1995: 30), Eriksson & Winka (1997: 4), etc, *nomina nuda*.

Basic type: *Peziza* Fr. 1822.

(Table 2, Fig 2) Cavalier-Smith was not the first to propose the phylum name *Ascomycota*. It appears to have been used first by Bold (1957: 7, 180), but without a Latin diagnosis. The name was in widespread use before its validation by Cavalier-Smith, and its usage was popularized by its employment in the eighth edition of the Dictionary, which is listed in Cavalier-Smith’s (1998) bibliography. The Latin diagnosis provided by Cavalier-Smith consisted of only two words: ‘sporaе intracellulares’. It is questionable whether this description is diagnostic for the *Ascomycota*, but as a validating diagnosis it is acceptable under the Code. No detailed reference to the basionym was given, but is provided here. We also propose a basic type, *Peziza*, as we can not be sure that the phylum will not be split in the future when more molecular data and material of ascomycetes and basidiomycetes have been sequenced. Hawksworth *et al.* (1995) and Eriksson & Winka (1997: 4) used the phylum names *Ascomycota* and *Basidiomycota*; the latter authors listed 31 nucleotide signatures in the nSSU rDNA genes in *Basidiomycota*. Since then many more sequences have become available, also from many other genes that support monophyly of *Ascomycota* and *Basidiomycota*.

The subdivision of *Ascomycota* used in the present paper is based on the system of Eriksson & Winka (1997), which differs in many respects from that of Cavalier-Smith (1998).

Subphylum: **Taphrinomycotina** O. E. Erikss. & Winka, *Myconet* 1: 11 (1997).

Class: **Taphrinomycetes** O. E. Erikss. & Winka, *Myconet* 1: 11 (1997).

Order: **Taphrinales** Gäum. & C. W. Dodge, *Comp. morph. fun.*: 159 (1928).

Exemplar genera: *Taphrina* Fr. 1815, *Protomyces* Unger 1832.

Class: **Neoelectomycetes** O. E. Erikss. & Winka, *Myconet* 1: 8 (1997).

Order: **Neoelectales** Landvik, O. E. Erikss, Gargas & P. Gustafss., *Syst. Ascom.* 11: 114 (1993).

Exemplar genus: *Neoelecta* Speg. 1881.

Class: **Pneumocystidomycetes** O. E. Erikss. & Winka, *Myconet* 1: 9 (1997).

Order: **Pneumocystidales** O. E. Erikss., *Syst. Ascom.* 13: 170 (1994).

Exemplar genus: *Pneumocystis* P. Delanoë & Delanoë 1912.

Class: **Schizosaccharomycetes** O. E. Erikss. & Winka, *Myconet* 1: 10 (1997).

Order: **Schizosaccharomycetales** O. E. Erikss., Svedskog & Landvik, *Syst. Ascom.* 11: 146 (1993).

Exemplar genus: *Schizosaccharomyces* Linder 1893.

Subphylum: **Saccharomycotina** O. E. Erikss. & Winka, *Myconet* 1: 10 (1997).

Class: **Saccharomycetes** O. E. Erikss. & Winka, *Myconet* 1: 10 (1997).

Order: **Saccharomycetales** Kudryavtsev, *System Hefen*: 270 (1960).

Growth usually by individual yeast cells, often accompanied by pseudohyphae and/or true hyphae. Cell walls predominately of β -glucan. Ascomata not formed; one to many ascospores formed in asci that often are converted from individual cells or borne on simple ascophores. Mitotic and meiotic nuclear divisions within an intact nuclear membrane. Enveloping membrane system in ascospore delimitation associated independently with postmeiotic nuclei. Asexual reproduction by holoblastic budding, conidia or fission (arthrospores).

Exemplar genera: *Saccharomyces* Meyen ex E. C. Hansen 1838, *Candida* Berkhout 1923, *Dipodascopsis* L. R. Batra & Millner 1978, *Metschnikowia* T. Kamieński 1899.

Subphylum: **Pezizomycotina** O. E. Erikss. & Winka, *Myconet* 1: 9 (1997).

Class: **Arthoniomycetes** O. E. Erikss. & Winka, *Myconet* 1: 4 (1997).

Order: **Arthoniales** Henssen & Jahns ex D. Hawksw. & O. E. Erikss, *Syst. Ascom.* 5: 177 (1986).

Synonym: *Arthoniales* Henssen & Jahns, *Lichenes*: 123 (1973) [‘1974’], *nomen invalidum*.

Hawksworth & Eriksson (*loc. cit.*) listed only Henssen, but cited the book by Henssen & Jahns (*loc. cit.*) as place for the original but invalid description so both should be cited although Henssen contributed the taxonomic system to the book.

Exemplar genera: *Arthonia* Ach. 1806, *Chrysothrix* Mont. 1852, *Dirina* Fr. 1825, *Roccella* DC. 1805.

Class: **Dothideomycetes** O. E. Erikss. & Winka, *Myconet* 1: 5 (1997).

Table 2 – Support for major groups of Fungi in selected phylogenetic studies: Ascomycota

Rank	Taxon	Reference	Data	OTUs	Support
Phylum	ASCOMYCOTA	James et al. (2006, fig. 1)	SSU, LSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	111	MLBS = 94 BPP = 1
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	177	WPBS = < 50 MLBS = 100 BPP = 1
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	276	NJBS = 67 BPP = 1
Subphylum	<i>Taphrinomycotina</i>	James et al. (2006, fig. 2)	SSU, LSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	4	MLBS = 98 BPP = 1
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	8	WPBS = < 50 MLBS = 98 BPP = 1
		Liu et al. (2006, fig. 3)	<i>rpb1</i> , <i>rpb2</i>	3	BPP = 1
		Sugiyama et al. (2007, fig. 2)	LSU, SSU <i>rpb2</i> , β - <i>tub</i>	11	BPP = 1
		Kurtzman & Sugiyama (2001, fig. 7)	SSU	8	NJBS = 54
Class/Order	<i>Taphrinomycetes</i> , <i>Taphrinales</i>	Sugiyama et al. (2007, fig. 2)	LSU, SSU <i>rpb2</i> , β - <i>tub</i>	6	BPP = 1
		Kurtzman & Sugiyama (2001, fig. 7)	SSU	4	NJBS = 100
Class/Order	<i>Neoelectomycetes</i> , <i>Neoelectales</i>	Nishida & Sugiyama (1994, fig. 1)	SSU	5	NJBS = 100
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	1	NA
Class/Order	<i>Pneumocystidomycetes</i> , <i>Pneumocystidales</i>	Sugiyama et al. (2007, fig. 2)	LSU, SSU, β - <i>tub</i> , <i>rpb2</i>	2	BPP = 1
		Landvik et al. (2001, fig. 1)	β - <i>tub</i>	2	MPBS = 100
Class/Order	<i>Schizosaccharomycetes</i> , <i>Schizosaccharomycetales</i>	Sugiyama et al. (2007, fig. 2)	LSU, SSU, β - <i>tub</i> , <i>rpb2</i>	1	NA
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	1	NA
Class/Order	<i>Schizosaccharomycetes</i> , <i>Schizosaccharomycetales</i>	Sugiyama et al. (2007, fig. 2)	SSU, LSU, <i>rpb2</i> , β - <i>tub</i>	1	NA
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	2	BPP = 1.0 NJBS = 100
Genus	<i>Taphrinomycotina incertae sedis</i> (not placed in any subphylum) <i>Saitoella</i>	Sugiyama et al. (2007, fig. 2)	SSU, LSU, <i>rpb2</i> , β - <i>tub</i>	1	NA
		Nishida & Sugiyama (1994, fig. 1)	SSU	1	NA
Subphylum/Class/Order	<i>Saccharomycotina</i> , <i>Saccharomycetes</i> , <i>Saccharomycetales</i>	Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	12	WPBS = 55 MLBS = 100 BPP = 1
		Suh et al. (2007, fig. 2)	LSU, SSU	87	MPBS = 99 BPP = 1
Subphylum	<i>Pezizomycotina</i>	James et al. (2006, fig. 1)	SSU, LSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	46	BPP = 1 MLBS = 94
		Robbertse et al. (2006, figs. 4,5,6)	Genomes	11	MPBS = 94-100 NJBS = 100 MLBS = 100
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	157	WPBS = 100 MLBS = 97 BPP = 1
Class/Order	<i>Arthoniomycetes</i> , <i>Arthoniales</i>	Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	4	WPBS = 100 MLBS = 100 BPP = 1
		Lumbsch et al. (2005, fig. 1)	LSU, SSU, mt-SSU, mt-LSU	6	MPBS = 100 BPP = 1.0
Class	<i>Dothideomycetes</i>	Schoch et al. (2007, fig.1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	96	BPP = 1 MPBS < 50 MLBS = 70
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	17	WPBS < 50 MLBS = 84 BPP = 1
		Kruys et al. (2006, fig. 1)	LSU, SSU, mt-SSU	51	BPP > 0.95 MPBS < 50
Subclass	<i>Dothideomycetidae</i>	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	26	BPP = 1 MPBS > 50 MLBS > 0.7

Table 2 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
		Kruys et al. (2006, fig. 1)	LSU, SSU, mt-SSU	11	BPP > 0.95 MPBS < 50
Order	Capnodiales	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	11	BPP = 1 MPBS > 70 MLBS > 70
Order	Dothideales	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	9	BPP = 1 MPBS > 70 MLBS > 70
		Kruys et al. (2006, fig. 1)	LSU, SSU, mt-SSU	4	BPP > 0.95 MPBS = 100
		Lindemuth et al. (2001)	LSU, SSU	6	MLBS = 91 NJBS = 100
Order	Myriangiales	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	5	BPP = 1 MPBS > 70 MLBS > 70
Subclass/Order	Pleosporomycetidae, Pleosporales	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	48	BPP = 1 MPBS > 70 MLBS > 70
		Kruys et al. (2006, fig. 1)	LSU, SSU, mt-SSU	35	BPP = 1 MPBS = 100
	<i>Dothideomycetes</i> <i>incertae sedis</i> (not placed in any subclass)				
Order	Botryosphaeriales	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	8	BPP = 1 MPBS > 70 MLBS > 70
Order	Hysteriales	Schoch et al. (2007, fig. 1)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	3	BPP = 1 MPBS > 70 MLBS > 70
Order	Patellariales	Pang et al. (2002, fig. 26)	SSU	1	NA
		Inderbitzin et al. (2001, fig. 18)	SSU	1	NA
Order	Jahnulales	Pang et al. (2002, fig. 26)	SSU	6	MPBS = 100
Class	Eurotiomycetes	Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	11	WPBS = 89 MLBS = 84
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	49	BPP = 1 MPBS = 100 WPBS = 100 MLBS = 100
		Ekman & Tønnsberg (2002, fig. 1)	SSU	13	BPP = 0.99
		Del Prado et al. (2006, fig. 1)	LSU, mt-SSU	15	BPP = 1
		Lumbsch et al. (2005, fig. 1)	LSU, SSU, mt-SSU, mt-LSU	11	BPP > 0.95 MPBS > 70
		Lutzoni et al. (2004, fig. 5)	LSU, SSU, mt-SSU, <i>rpb2</i>	8	BPP = 1 BBS = 61
		Reeb et al. (2004, fig. 1)	SSU, LSU, <i>rpb2</i>	7	BPP = 1 BBS = 89
Subclass	Chaetothyriomycetidae	Reeb et al. (2004, fig. 1)	SSU, LSU, <i>rpb2</i>	5	BPP = 1 BBS = 100 MLBS = 100
		Lutzoni et al. (2004, fig. 5)	LSU, SSU, mt-SSUSSU, <i>rpb2</i>	5	BPP = 1 BBS = 100 NJBS = 99 MPBS = 98
		Del Prado et al. (2006, fig. 1)	LSU, mt-SSU	11	BPP = 1
		Spatafora et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	6	BPP = 1 MLBS = 100 WPBS > 70
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	21	BPP = 1 MPBS = 100 WPBS = 100 MLBS = 100
Order	Chaetothyriales	Lutzoni et al. (2004, fig. 2)	LSU, SSU	5	BPP = 1 NJBS = 94

(continued on next page)

Table 2 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
Order	Pyrenulales	Liu & Hall (2004, fig. 3)	<i>rpb2</i>	5	BPP = 1 MPBS = 96
		Spatafora et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	4	BPP = 1 MLBS = 100 WPBS > 70
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	9	BPP = 1 MPBS = 100 WPBBS = 100 MLBS = 100
		Lutzoni et al. (2004, fig. 8)	LSU, SSU, mt-SSU, <i>rpb2</i>	2	BPP = 1 NJBS = 100 WPBS = 100
		Reeb et al. (2004, fig. 1)	LSU, SSU, <i>rpb2</i>	2	BPP = 1 BBS = 100 MLBS = 100
Order	Verrucariales	Schmitt et al. (2004, fig. 1)	LSU, mt-SSU	2	BPP = 1
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	5	BPP = 1 MPBS = 100 WPBBS = 100 MLBS = 100
		Wedin et al. (2006, fig. 1)	LSU, mt-SSU	3	BPP = 1 MPjk = 100
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	7	BPP = 1 MPBS = 100 WPBBS = 100 MLBS = 100
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	3	BPP = 1 NJBS = 98
Subclass	Eurotiomycetidae	Gueidan et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i>	83	BPP = 1 MLBS = 100 MPBS = 100
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	24	BPP = 1 MPBS = 100 WPBBS = 98 MLBS = 100
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	11	NJBS = 96 BPP = 1
Order	Coryneliales	Winka (2000, fig. 1)	SSU	2	MPBS = 100 NJBS = 100
		Inderbitzin et al. (2004, fig. 14)	SSU	1	NA
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	3	BPP = 1 MPBS = 100 WPBBS = 100 MLBS = 100
Order	Eurotiales	Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	9	BPP = 1 MPBS = 100 WPBBS = 100 MLBS = 100
Order	Onygenales	Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	12	BPP = 1 MPBS = 65 WPBBS = 68 MLBS = 88
Subclass/Order	Mycocaliciomycetidae, Mycocaliciales	Tibell & Vinuesa (2005, fig. 1)	LSU	20	BPP = 1
		Geiser et al. (2007, fig. 1)	SSU, LSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef</i>	4	BPP = 1 MPBS = 100 WPBBS = 100 MLBS = 100
Class	Laboulbeniomycetes	Ekman & Tønberg (2002, fig. 1)	SSU	4	BPP = 1
Order	Laboulbeniales	Weir & Blackwell (2001, fig. 2)	SSU	4	MPBS = 100
		Henk et al. (2003, fig. 1)	SSU	6	MPBS = 100
		Weir & Blackwell (2001, fig. 1)	SSU	3	MPBS = 100
Order	Pyxidiphorales	Henk et al. (2003, fig. 2)	SSU	3	MPBS = 57
		Weir & Blackwell (2001, fig. 2)	SSU	1	NA
Order		Henk et al. (2003, fig. 2)	SSU	2	MPBS = 99

Table 2 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
Class	<i>Lecanoromycetes</i>	Lutzoni et al. (2004, fig. 5)	LSU, SSU, <i>rpb2</i> , mt-SSU	34	BPP = 1 BBS = 56
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	38	WPBS < 50 MLBS = 93 BPP = 1
		Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	264	RMLBS > 70 BS BPP > 0.95
		Hofstetter et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	82	RMLBS > 70 BPP > 0.95
Subclass/Order	<i>Acarosporomycetidae</i> , <i>Acarosporales</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	15	RMLBS > 70 % PMLBS > 70 % BPP > 0.95
		Reeb et al. (2004, fig. 1)	LSU, SSU, <i>rpb2</i>	14	MLBS = 100 BPP = 100
		Lutzoni et al. (2004, fig. 4)	LSU, SSU, <i>rpb2</i>	14	BPP = 1 NJBS = 100 MPBS = 100
Subclass	<i>Lecanoromycetidae</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	71	RMLBS > 70 % PMLBS > 70 % BPP > 0.95
		Hofstetter et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	54	RMLBS > 70 BPP > 0.95
		Reeb et al. (2004, fig. 1)	LSU, SSU, <i>rpb2</i>	14	MLBS = 73 BPP = 100
Order	<i>Lecanorales</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	86	RMLBS > 70 BS BPP > 0.95
		Hofstetter et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	30	RMLBS > 70 BPP > 0.95
		Lumbsch et al. (2004, fig. 1) Lücking et al. (2004, fig. 3)	LSU, mt-SSU LSU, mt-SSU	14 8	BPP = 1 BPP = 1
Order	<i>Peltigerales</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	46	RMLBS > 70 BSBPP > 0.95
		Miądlikowska & Lutzoni (2004, fig. 1)	LSU, SSU	59	MPBS < 70 BPP = 0.92
Order	<i>Teloschistales</i>	Wilklund & Wedin (2003, fig. 1)	LSU, SSU	31	Bjk = 99
		Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	13	RMLBS > 70 BPP > 0.95
Subclass	<i>Ostropomycetidae</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	58	RMLBS > 70 BS BPP > 0.95
		Grube et al. (2004, fig. 1) Reeb et al. (2004, fig. 1)	mt-SSU LSU, SSU, <i>rpb2</i>	30 16	BPP > 0.95 MLBS = 100 BPP = 100
Order	<i>Agyriales</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	8	RMLBS > 70 BS BPP > 0.95
		Lücking et al. (2004, fig. 3) Lutzoni et al. (2004, fig. 2)	LSU, mt-SSU LSU, SSU	11 4	BPP = 1 BPP = 1 NJBS = 100
		Wedin et al. (2005, fig. 1)	LSU, mt-SSU	8	MPjk = 83 BPP = 0.99
Order	<i>Baeomycetales</i>	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	4	RMLBS > 70 PMLBS > 70 BPP > 0.95
		Wedin et al. (2005, fig. 1)	LSU, mt-SSU	3	MPjk = 99 BPP = 1.0
Order	<i>Ostropales</i> s.l.	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	21	RMLBS > 70 BS BPP > 0.95
		Schmitt et al. (2005, fig. 1)	LSU, mt-SSU	12	BPP = 1
		Wedin et al. (2005, fig. 1)	LSU, mt-SSU	13	Bjk = 94 BPP = 0.97

(continued on next page)

Table 2 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
Order	Pertusariales	Lutzoni et al. (2004, fig. 4)	LSU, SSU, <i>rpb2</i>	10	BPP = 1 NJBS = 74 MPBS = 84
		Reeb et al. (2004, fig. 1)	LSU, SSU, <i>rpb2</i>	9	MLBS = 99 BPP = 1 BBS = 1
		Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	21	RMLBS > 70 BSBpp > 0.95
		Lücking et al. (2004, fig. 3)	LSU, mt-SSU	7	BPP = 1
		Schmitt et al. (2005, fig. 1)	LSU, mt-SSU	14	BPP = 1
Order	Lecanoromycetes incertae sedis (not placed in any subclass) Candelariales	Lutzoni et al. (2004, fig. 2)	LSU, SSU	11	BPP = 1
		Wedin et al. (2005, fig. 1)	LSU, mt-SSU	3	Jk = 100 BPP = 0.96
		Hofstetter et al. (2007, fig. 1)	LSU, SSU, mt-SSU, <i>rpb1</i> , <i>rpb2</i>	2	RMLBS > 70 BPP > 0.95
		Miądlikowska et al. (2007, fig. 1)	LSU, SSU, mt-SSU, <i>rpb1</i> , <i>rpb2</i>	3	RMLBS > 70 PMLBS > 70 BPP > 0.95
Order	Umbilicariales	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	16	BSBSBPP > 0.95
		Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	9	RMLBS > 70 PMLBS > 70 BPP > 0.95
		Hofstetter et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	8	RMLBS > 70 BPP > 0.95
		Reeb et al. (2004, fig. 1)	LSU, SSU, <i>rpb2</i>	4	MLBS = 70 BPP = 1 BBS = 88
Class	Leotiomycetes (w/o Geoglossaceae)	Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	22	WPBS = 100 MLBS = 100 BPP = 1
		Wang et al. (2006, fig. 1)	LSU, SSU, 5.8S	50	BPP = 1
		Wang et al. (2007, fig. 2)	LSU, SSU, 5.8S	78	MPBS = 61 BPP = 1
Order	Cyttariales	Wang et al. (2007, fig. 1)	SSU, LSU, 5.8S	1	NA
Order	Erysiphales	Rossmann et al. (2004, fig. 2)	LSU	12	MPBS > 55
		Wang et al. (2007, fig. 1)	SSU, LSU, 5.8S	16	MPBS = 63 BPP = 0.97
Order	Helotiales (w/o Geoglossaceae)	Takamatsu (2004, fig. 2)	SSU	10	NJBS = 99
		Wang et al. (2007, fig. 1)	SSU, LSU, 5.8S	40	BPP < 0.90
Order	Rhytismatales	Rossmann et al. (2004, fig. 2)	LSU	4	MPBS > 55
		Wang et al. (2007, fig. 1)	SSU, LSU, 5.8S	5	MPBS = 100 BPP = 1
Order Class/Order	Thelebolales Lichinomycetes, Lichinales	de Hoog et al. (2005, fig. 3)	SSU	11	MPBS = 56
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	1	NA
Class/Order	Orbiliomycetes, Orbiliales	Miądlikowska et al. (2007, fig. 1)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , mt-SSU	2	RMLBS > 70 PMLBS > 70 BPP > 0.95
		Reeb et al. (2004, fig. 1)	LSU, SSU, <i>rpb2</i>	3	MLBS = 100 BBS = 100 BPP = 1
		Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	2	WPBS = 100 MLBS = 100 BPP = 1
Class/Order	Pezizomycetes, Pezizales	Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	14	WPBS = 54 MLBS = 99 BPP = 1
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	21	BPP = 0.96 NJBS = 70
Class	Sordariomycetes	Spatafora et al. (2007, fig. 2)	LSU, SSU, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	47	WPBS = 100 MLBS = 100 BPP = 1

Table 2 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
		Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	106	MPBS = 100 WPBS = 100 MLBS = 100 BPP = 1
Subclass	<i>Hypocreomycetidae</i>	Lutzoni et al. (2004, fig. 2)	LSU, SSU	66	BPP = 1 NJBS = 97
		Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	42	MPBS = 92 WPBS = 96 MLBS = 90 BPP = 1
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	26	NJBS < 50 BPP = 1
Order	<i>Coronophorales</i>	Huhndorf et al. (2004b, figs. 38, 39)	LSU	21	MPBS = 67 BPP >= 0.95
		Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	2	MPBS < 50 WPBS < 50 MLBS = 96 BPP = 1
		Huhndorf et al. (2004b, figs. 38, 39)	LSU	16	WPBS = 99 BPP ≥ 95
Order	<i>Hypocreales</i>	Miller & Huhndorf (2005, fig. 7)	LSU, β - <i>tub</i> , <i>rpb2</i>	2	WPBS = 100 BPP ≥ 95
		Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	21	MPBS = 91 WPBS = 90 MLBS = 72 BPP = 1
		Castlebury et al. (2004, fig. 1)	LSU, SSU	31	MPBS = 70 BPP = 1
Order	<i>Melanosporales</i>	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	2	MPBS = 100 WPBS = 100 MLBS = 100 BPP = 1
Order	<i>Microascales</i> (incl. <i>Halosphaeriales</i>)	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	15	MPBS = 74 WPBS = 86 MLBS = 85 BPP = 1
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	10	NJBS = 80 BPP = 1
		Campbell et al. (2003, fig. 3)	LSU, SSU	40	MPBS = 100 BPP = 1
Subclass	<i>Sordariomycetidae</i>	Kohlmeyer et al. (2000, fig. 1)	LSU, SSU	16	MPBS = 97
		Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	54	MPBS = 82 WPBS = 85 MLBS = 77 BPP = 1
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	36	NJBS < 50 BPP = 0.97
Order	<i>Boliniales</i>	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	4	MPBS = 100 WPBS = 100 MLBS = 100 BPP = 1
		Huhndorf et al. (2004a, fig. 1)	LSU	3	WPBS = 99 BPP < 95
		Miller & Huhndorf (2005, fig. 7)	LSU, β - <i>tub</i> , <i>rpb2</i>	2	WPBS = 100 BPP ≥ 95
Order	<i>Chaetosphaeriales</i>	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	3	MPBS = 100 WPBS = 100 MLBS = 100 BPP = 1
		Miller & Huhndorf (2005, fig. 7)	LSU, β - <i>tub</i> , <i>rpb2</i>	2	WPBS = 100 BPP ≥ 95
		Shenoy et al. (2006, fig. 3)	LSU, <i>rpb2</i>	4	MPBS = 100

(continued on next page)

Table 2 (continued)

Rank	Taxon	Reference	Data	OTUs	Support	
Order	Coniochaetales	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	3	MPBS = 93 WPBS = 100 MLBS = 87 BPP = 1	
		Miller & Huhndorf (2005, fig. 7)	LSU, β - <i>tub</i> , <i>rpb2</i>	2	WPBS = 100 BPP \geq 95	
		Miller & Huhndorf (2004, fig. 10)	LSU	3	WPBS = 98 BPP \geq 95	
Order	Diaporthales	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	19	MPBS = 95 WPBS = 94 MLBS = 77 BPP = 1	
		Castlebury et al. (2002, fig. 1)	LSU	82	MPBS = 100 NJBS = 100	
		Lutzoni et al. (2004, fig. 2)	LSU, SSU	10	NJBS = 100 BPP = 1	
		Miller & Huhndorf (2005, fig. 7)	LSU, β - <i>tub</i> , <i>rpb2</i>	2	WPBS = 100 BPP \geq 95	
		Miller & Huhndorf (2004, fig. 10)	LSU	3	WPBS = 100 BPP \geq 95	
Order	Ophiostomatales	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	3	MPBS = 100 WPBS = 100 MLBS = 100 BPP = 1	
		Hausner & Reid (2004, fig. 1) Wingfield et al. (1999, fig. 3)	SSU LSU	3 4	NJBS = 99 MPBS = 99	
Order	Sordariales	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	17	MPBS = 80 WPBS = 77 MLBS = 84 BPP = 1	
		Huhndorf et al. (2004a, fig. 1)	LSU	22	WPBS = < 50 BPP < 95	
		Miller & Huhndorf (2005, fig. 7)	LSU, β - <i>tub</i> , <i>rpb2</i>	41	WPBS = 65 BPP \geq 95	
		Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	8	MPBS = 98 WPBS = 99 MLBS = 78 BPP = 1	
Subclass/Order	Xylariomycetidae, Xylariales	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	16	MPBS = 92	
Order	Sordariomycetes incertae sedis (not placed in any subclass)	Shenoy et al. (2006, fig. 1)	LSU	16	MPBS = 92	
		Calosphaeriales	Vijaykrishna et al. (2004, fig. 1)	SSU	3	MPBS = 100
			Réblová et al. (2004, fig. 1) Réblová (2006, fig. 1)	LSU SSU	6 2	MPBS = 53 MPBS = 68
Order	Lulworthiales (incl. <i>Spathulosporales</i>)	Zhang et al. (2007, fig. 2)	LSU, SSU, <i>rpb2</i> , <i>tef1</i>	2	MPBS = 100 WPBS = 100 MLBS = 100 BPP = 1	
		Campbell et al. (2005, fig. 1)	LSU, SSU	56	BPP = 1	
		Inderbitzin et al. (2004, fig. 15)	LSU	15	MPBS = 100 NJBS = 91 BPP = 86	
		Kohlmeyer et al. (2000, fig. 1)	LSU, SSU	7	MPBS = 100	
Order	Meliolales	Saenz & Taylor (1999, fig. 1)	LSU	2	MPBS = 100	
Order	Phyllachorales	Vijaykrishna et al. (2004, fig. 1)	SSU	2	MPBS < 50	
Order	Trichosphaeriales	Inderbitzin et al. (2004, fig. 14)	SSU	1	NA	
Order	Pezizomycotina incertae sedis (not placed in any class)	Réblová & Seifert (2004, fig. 1)	LSU	8	MPBS < 50	
Order	Lahmiales	Eriksson (1986)	—	—	—	
Order	Medeolariales	Inderbitzin et al. (2004, fig. 14)	SSU	1	NA	
Order	Triblidiales	Eriksson (1992)	—	—	—	

See Table 1 for explanation.




Fig 2 – Phylogeny and classification of Fungi. Ascomycota. See Table 2 for support values for clades. Dashed lines indicate taxa that are of uncertain placement.

Subclass: **Dothideomycetidae** P. M. Kirk, P. F. Cannon, J. C. David & Stalpers ex Schoch et al., *Mycologia* 98: 1047 (2007) ['2007'].

Order: **Capnodiales** Woron., *Annls Mycol.* 23: 177 (1925).
Exemplar genera: *Capnodium* Mont. 1848, *Scorias* Fr. 1825, *Mycosphaerella* Johanson 1884.

Order: **Dothideales** Lindau, in Engler & Prantl (eds), *Nat. Pflanzenfam.* 1(1): 373 (1897).
Exemplar genera: *Dothidea* Fr. 1818, *Dothiora* Fr. 1849, *Sydowia* Bres. 1895, *Stylodothis* Arx & E. Müll. 1975.

Order: **Myriangiales** Starbäck, *K. svenska Vetensk-Akad. Handl., Bih., Afd. III* 25: 37 (1899).
Exemplar genera: *Myriangium* Mont. & Berk. 1845, *Elsinoë* Racib. 1900.

Subclass: **Pleosporomycetidae** C. L. Schoch, Spatafora, Crous & Shoemaker, *Mycologia* 98: 1049 (2007) ['2006'].

Order: **Pleosporales** Luttr. ex M. E. Barr, *Prodr. Class Loculoasc.* 67 (1987b).

Synonym: **Pleosporales** Luttr., *Mycologia* 47: 520 (1955), nomen invalidum.

Exemplar genera: *Pleospora* Rabenh. ex Ces. & De Not. 1863, *Phaeosphaeria* I. Miyake 1909, *Lophiostoma* Ces. & De Not. 1863, *Sporormiella* Ellis & Everh. 1892, *Montagnula* Berl. 1896.

Dothideomycetes incertae sedis (not placed in any subclass)

Order: **Botryosphaerales** C. L. Schoch, Crous & Shoemaker, *Mycologia* 98: 1051 (2007) ['2006'].

Exemplar genera: *Botryosphaeria* Ces. & De Not. 1863, *Guignardia* Viala & Ravaz 1892.

Order: **Hysteriales** Lindau in Engler & Prantl (eds), *Nat. Pflanzenfam.* 1: 265 (1896), as 'Hysteriinae'.

Exemplar genera: *Hysterium* Pers. 1797, *Hysteropatella* Rehm. 1890.

Order: **Patellariales** D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 181 (1986).

Exemplar genus: *Patellaria* Fr. 1822.

Order: **Jahnulales** Ka-Lai Pang, Abdel-Wahab, El-Shar., E. B. G. Jones & Sivichai, in Pang et al., *Mycol. Res.* 106: 1033 (2002).

Exemplar genera: *Aliquandostipite* Inderb. 2001, *Jahnula* Kirschst. 1936, *Patescospora* Abdel-Wahab & El-Shar. 2002.

Class: **Eurotiomycetes** O. E. Erikss. & Winka, *Myconet* 1: 6 (1997).

The circumscription of this class and the classification within the **Eurotiomycetes** presented here are derived from the phylogenetic re-delimitation of this class by Ekman & Tønberg (2002), Lutzoni et al. (2004) and Geiser et al. (2007), reflecting the inference of shared ancestry between **Eurotiomycetes**, comprising **Coryneliales**, **Onygenales** and **Eurotiales** and **Chaetothyriomycetes**. Three subclasses, **Chaetothyriomycetidae**, **Eurotiomycetidae**, and **Mycocaliciomycetidae**, are defined to represent the major lineages within **Eurotiomycetes**.

Subclass: **Chaetothyriomycetidae** Doweld, *Prosyllabus*: LXXVIII (2001).

Lichenized, parasitic, and saprobic ascomycetes with mostly bitunicate/fissitunicate to evanescent asci, produced in perithecial ascomata arranged superficially or immersed in a thallus. Thalli often produced on the surfaces of rocks, lichens, decaying plant material and other substrata. Ascospores variable, from colourless to pigmented, simple to muriform. Hamathecium, when present, consisting of pseudoparaphyses. Pigments, when present, generally related to melanin. Asexual stages with phialidic and anellidic anamorphs observed in non-lichenized taxa.

Order: **Chaetothyriales** M. E. Barr, *Mycotaxon* 29: 502 (1987).

Exemplar genera: *Capronia* Sacc. 1883, *Ceramothyrium* Bat. & H. Maia 1956, *Chaetothyrium* Speg. 1888.

Order: **Pyrenulales** Fink ex D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 182 (1986).

Synonym: *Pyrenulales* Fink, *Ohio St. Univ. Bull.* 19(28): 107 (1951), *nomen invalidum*.

Exemplar genera: *Pyrenula* Ach. 1814, *Pyrgillus* Nyl. 1858.

Order: **Verrucariales** Mattick ex D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 183 (1986).

Synonym: *Verrucariales* Mattick, in Engler, *Syll. Pflanzenfam.* (12 edn): 208 (1954), *nomen invalidum*.

Exemplar genera: *Agonimia* Zahlbr. 1909, *Dermatocarpon* Eschw. 1824, *Polyblastia* A. Massal. 1852, *Verrucaria* Schrad. 1794.

Subclass: **Eurotiomycetidae** Geiser & Lutzoni, **subclass. nov.**
Mycobank no.: MB 501287

Fungi saprotrophici vel parasitici vel mycorrhizales; asci globosi in toto ascomate sparsi, raro hymenium formantes; asci plerumque evanescentes, nonnumquam bitunicati. Ascosporeae plerumque unicellulares, lenticulares, nonnumquam globosae vel ellipsoideae. Ascomata, si formata, plerumque cleistothecialia vel gymnothecialia, saepe textura stromatica circumdata. Structurae hamatheciales absentes. Gametangia plerumque indistincta e glomere hyphali constantia. Fungi saepe laete colorati. Anamorphae variables, seu phialidicae seu arthroconidiales.

Typus: *Eurotium* Link 1809.

Saprotrophic, parasitic and mycorrhizal. Ascomata, when present, usually cleistothecial/gymnothecial, globose, often produced in surrounding stromatic tissue and brightly coloured; hamathecial elements lacking; gametangia usually undifferentiated and consisting of hyphal coils. Asci usually evanescent, sometimes bitunicate, scattered throughout the ascoma, rarely from a hymenium. Ascospores usually single-celled, lenticular, sometimes spherical or elliptical. Anamorphs variable, including phialidic and arthroconidial forms.

This name was employed by Lutzoni et al. (2004) and Geiser et al. (2007), in the same sense as the present classification, but without a formal diagnosis.

Order: **Coryneliales** Seaver & Chardón, *Scient. Surv. P. Rico*: 40 (1926).

Exemplar genera: *Corynelia* Ach. 1823, *Caliciopsis* Peck 1880.

Order: **Eurotiales** G. W. Martin ex Benny & Kimbr., *Mycotaxon* 12: 23 (1980).

Synonym: *Eurotiales* G. W. Martin, *Std. nat. Hist. Iowa Univ.* 18(Suppl.): 16 (1941), *nomen invalidum*.

Exemplar genera: *Eurotium* Link 1809, *Emericella* Berk. 1857, *Talaromyces* C. R. Benj. 1955, *Elaphomyces* Nees 1820, *Trichocoma* Jungh. 1838, *Byssoschlamys* Westling 1909.

Order: **Onygenales** Cif. ex Benny & Kimbr., *Mycotaxon* 12: 8 (1980).
Synonym: *Onygenales* Cif., *Atti Ist. Bot. Univ. Pavia, ser. 5*, 14: 238 (1957), *nomen invalidum*.

Emend. Currah *Mycotaxon* 24: 13 (1985).

Exemplar genera: *Onygena* Pers. 1799, *Gymnoascus* Baran. 1872, *Arthroderma* Curr. 1860.

Subclass: **Mycocaliciomycetidae** Tibell. **subclass nov.**
Mycobank no.: MB 501288

Parasitae vel commensales in lichenibus vel saprotrophici. Ascomata disciformia, stipitata vel sessilia. Excipulum cupulatum, saltem partim scleroticum hyphis stipitis simile. Dispersio sporarum activa, raro passiva et tum mazedio parce evoluto. Asci unitunicati, cylindrici, vulgo apice distincte incrassato, 8-spori. Ascosporeae pallidae ad atrofuscae, ellipsoideae, non-septatae vel transversaliter 1–7-septatae. Paries sporeae atrofuscae, laevis vel ornamento intra plasmalemma formato. Derivata acidi vulpinici in speciebus paucis praesentia. Anamorphae coelomycetum et hyphomycetum variae praesentes.

Typus: *Mycocalicium* Vain. 1890.

Parasites or commensals on lichens or saprobes. Ascomata disciform, stalked or sessile. Excipulum cupulate, and like the stalk hyphae at least in part sclerotized. Spore dispersal active, more rarely passive and ascomata then with a moderately developed mazaedium. Asci unitunicate, cylindrical, mostly with a distinctly thickened apex, 8-spored. Ascospores pale to blackish brown, ellipsoidal or spherical to cuboid, non-septate or transversely 1–7-septate. Spore wall pigmented, smooth or with an ornamentation formed within the plasmalemma. Vulpinic acid derivatives occur in a few species. A variety of coelomycetous and hyphomycetous anamorphs occur.

Order: **Mycocaliciales** Tibell & Wedin, *Mycologia* 92: 579 (2000).

Exemplar genera: *Mycocalicium* Vain. 1890, *Chaenothecopsis* Vain. 1927, *Stenocybe* (Nyl.) Körb. 1855, *Sphinctrina* Fr. 1825.

Class: **Laboulbeniomycetes** Engl., *Syll. Pflanzenfam.* (2nd edn): 46 (1898).

Order: **Laboulbeniales** Lindau, in Engler & Prantl (eds), *Nat. Pflanzenfam.* 1(1): 491 (1897), as ‘*Laboulbeniineae*’.

Exemplar genera: *Laboulbenia* Mont. & C.P. Robin 1835, *Rickia* Cavara 1899, *Ceratomyces* Thaxt. 1892.

Order: **Pyxidiophorales** P. F. Cannon, in Kirk et al., *Ainsworth & Bisby's Dict. Fungi* (9th edn): xi (2001).

Exemplar genus: *Pyxidiophora* Bref. & Tavel 1891.

Class: **Lecanoromycetes** O. E. Erikss. & Winka, *Myconet* 1: 7 (1997).

Subclass: **Acarosporomycetidae** Reeb, Lutzoni & Cl. Roux, *Mol. Phylogen. Evol.* 32: 1053 (2004).

Order: **Acarosporales** Reeb, Lutzoni & Cl. Roux, **ord. nov.**
Mycobank no.: MB 501289

Ascomycetes lichenisati algas virides thallo continentes. Ascomata immersa vel sessilia, disciformia vel peritheciodea. Excipulum hyalinum, annulatum. Hymenium non-amyloideum. Paraphyses mediocriter vel infirme ramosae, septatae, mediocriter vel infirme anastomosantes. Asci unitunicati, non-amyloidei vel satis infirme amyloidei, polyspori. Ascosporeae hyalinae, non-septatae, non-halonatae.

Typus: *Acarospora* A. Massal. 1852.

Lichen-forming ascomycetes with chlorococcoid photobiont. Ascomata immersed or sessile, disciform or perithecioid. True exciple hyaline, annulate. Hymenium non-

amyloid. Paraphyses moderately to poorly branched, septate, moderately to poorly anastomosing. Asci functionally unitunicate, lecanoralean, non-amyloid or with slightly amyloid tholi, polyspored, generally with more than 100 ascospores per ascus. Ascospores hyaline, small, non-septate, non-halonate.

The members of this order were formerly classified within the *Lecanorales*, but Reeb *et al.* (2004) and Lutzoni *et al.* (2004) demonstrated that the *Acarosporaceae* diverged earlier than the *Lecanoromycetidae* and *Ostropomycetidae*. This early divergence within the *Lecanoromycetes* was confirmed by Wedin *et al.* (2005) and Miądlikowska *et al.* (2007).

Exemplar genera: *Acarospora* A. Massal. 1852, *Pleopsidium* Körb. 1855, *Sarcogyne* Flot. 1851.

Subclass: **Lecanoromycetidae** P. M. Kirk, P. F. Cannon, J. C. David & Stalpers ex Miądlikowska, Lutzoni & Lumbsch, **subclass. nov.** MycoBank no.: MB 501290

Synonym: *Lecanoromycetidae* P. M. Kirk, P. F. Cannon, J. C. David & Stalpers, *Ainsworth & Bisby's Dict. Fungi* (9th edn): xi (2001), *nomen invalidum*.

Ascomycetes lichenisati algal virides vel cyanobacteria thallo continentis. Ascomata immersa, sessilia vel elevata, generaliter disciformia. Excipulum hyalinum vel pigmentatum, annulatum vel cupulatum. Hymenium amyloideum vel non-amyloideum. Paraphyses simplices vel ramosae, septatae, anastomosantes vel non-anastomosantes. Asci bitunicati, unitunicati vel prototunicati, non-amyloidei vel amyloidei, generaliter octospori, sed etiam 1- ad multispori. Ascosporae hyalinae vel brunneae, non-septatae, vel septatae usque ad muriformes, halonatae vel non-halonatae.

Typus: *Lecanora* Ach. 1809.

Lichen-forming ascomycetes with green algal or cyanobacterial photobiont. Ascomata immersed, sessile or stalked, usually disciform. True exciple hyaline or pigmented, annulate or cupulate. Hymenium amyloid or non-amyloid. Paraphyses simple or moderately to richly branched, septate, anastomosing or not. Asci bitunicate, functionally unitunicate, or prototunicate, lecanoralean, non-amyloid or amyloid, mostly 8-spored, but varying from 1- to poly-spored. Ascospores hyaline or brown, non-septate, trans-septate or muriform, halonate or non-halonate.

This subclass includes the bulk of lichenized discomycetes and corresponds to the phylogenetic circumscription of this subclass by Reeb *et al.* (2004), Lutzoni *et al.* (2004) and Miądlikowska *et al.* (2007). It is in agreement with the *Lecanorales* of Lumbsch *et al.* (2004) and Wiklund & Wedin (2004). The orders *Peltigerales* and *Teloschistales* are here accepted at the ordinal level, following Miądlikowska & Lutzoni (2003) and Miądlikowska *et al.* (2007).

Order: **Lecanorales** Nannf., *Nova Acta R. Soc. Scient. Upsal*, ser. 4 8(2): 68 (1932).

Exemplar genera: *Cladonia* Hill. ex P. Browne 1756, *Lecanora* Ach. 1809, *Parmelia* Ach. 1803, *Ramalina* Ach. 1809, *Usnea* Dill. ex Adans. 1763

Order: **Peltigerales** Walt. Watson, *New Phytologist* 28: 9 (1929).

Exemplar genera: *Coccocarpia* Pers. 1827, *Collema* F. H. Wigg. 1780, *Nephroma* Ach. 1810, *Pannaria* Del. ex Bory 1828, *Peltigera* Willd. 1787.

Order: **Teloschistales** D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 183 (1986).

Exemplar genera: *Caloplaca* Th. Fr. 1861, *Teloschistes* Norman 1853, *Xanthoria* (Fr.) Th. Fr. 1860.

Subclass: **Ostropomycetidae** Reeb, Lutzoni & Cl. Roux, *Mol. Phylogen. Evol.* 32: 1055 (2004).

Order: **Agyriales** Clem. & Shear, *Gen. Fungi*: 141 (1931).

Exemplar genera: *Agyrium* Fr. 1822, *Placopsis* (Nyl.) Linds. 1867, *Trapelia* M. Choisy 1929, *Trapeliopsis* Hertel & Gotth. Schneid. 1980.

Order: **Baeomycetales** Lumbsch, Huhndorf & Lutzoni, **ord. nov.**

MycoBank no.: MB 501291

Ascomycetes lichenisati algal virides thallo continentis. Ascomata elevata vel raro sessilia, disciformia. Excipulum hyalinum vel pigmentatum, annulatum vel cupulatum. Hymenium non-amyloideum. Paraphyses ramosae, septatae. Asci unitunicati, non-amyloidei vel satis infirme amyloidei, octospori. Ascosporae hyalinae, non-septatae vel septatae, halonatae vel non-halonatae.

Typus: *Baeomyces* Pers. 1794.

Lichen-forming ascomycetes with chlorococcoid photobiont. Ascomata sessile or rarely stalked, disciform. True exciple hyaline or pigmented, annulate or cupulate. Hymenium non-amyloid. Paraphyses moderately to richly branched, septate. Asci unitunicate, non-amyloid or with slightly amyloid tholi, 8-spored. Ascospores hyaline, non-septate or trans-septate, halonate or non-halonate.

Baeomycetales was shown to differ from *Agyriales* by Kauff & Lutzoni (2002) and this was confirmed by Miądlikowska *et al.* (2007) and Lumbsch *et al.* (2007).

Exemplar genera: *Ainoa* Lumbsch & I. Schmitt 2001, *Baeomyces* Pers. 1794, *Phyllobaeis* Gierl & Kalb 1993.

Order: **Ostropales** Nannf., *Nova Acta R. Soc. Scient. Upsal*, ser. 4 8(2): 68 (1932).

This order includes also taxa formerly classified in separate orders, such as *Gomphillales*, *Graphidales*, *Gyalectales* and *Trichotheliales*.

Exemplar genera: *Ostropa* Fr. 1825, *Stictis* Pers. 1799, *Gyalecta* Ach. 1808, *Gomphillus* Nyl. 1855, *Graphis* Adans. 1763, *Odontotrema* Nyl. 1858, *Porina* Müll. Arg. 1883, *Thelotrema* Ach. 1803.

Order: **Pertusariales** M. Choisy ex D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 181 (1986).

Synonym: *Pertusariales* M. Choisy, *Bull. mens. Soc. linn. Lyon* 18: 12 (1949), *nomen invalidum*.

This order may not be monophyletic as currently circumscribed, with *Ochrolechiaceae* and some groups of the heterogeneous *Pertusaria* clustering in a separate clade, but without support. Nonetheless, a cluster of taxa in a 'core' group of *Pertusariales* has been strongly supported as monophyletic in phylogenetic analyses by Miądlikowska *et al.* (2007), Lücking

et al. (2004), Schmitt et al. (2005), Lutzoni et al. (2004), and Grube et al. (2004).

Exemplar genera: *Coccotrema* Müll. Arg. 1888, *Icmadophila* Trevis. 1853, *Ochrolechia* A. Massal. 1852, *Pertusaria* DC. 1805.

Lecanoromycetes incertae sedis (not placed in any subclass):

Order: **Candelariales** Miadl., Lutzoni & Lumbsch, **ord. nov.**

MycoBank no.: MB 501292

Ascomycetes lichenisati algas virides thallo continentis. Ascumata sessilia, disciformia. Excipulum hyalinum, annulatum. Hymenium amyloideum. Paraphyses ramosae, septatae. Asci unitunicati, amyloidei, ad typum *Candelariae* dictum pertinentes, octo- vel saepe multisporei. Ascosporeae hyalinae, non-septatae vel raro 1-septatae.

Typus: *Candelaria* A. Massal. 1853.

Lichen-forming ascomycetes with chlorococcoid photobiont, predominantly nitrophilous. Thallus of various morphology, yellow to orange (pulvinic acid derivatives). Ascumata apothecial, sessile, with or without a distinct margin, yellow to orange. The ascumatal wall formed from densely septate twisted hyphae. paraphyses mostly simple. Excipulum hyaline, hymenium amyloid. Asci unitunicate of *Candelaria*-type with the amyloid lower part of the apical dome and broad apical cushion, often multispored. Ascospores hyaline, aseptate, rarely 1-septate.

Candelariales was shown to differ from *Lecanorales* by Wedin et al. (2005) and this was confirmed by Hofstetter et al. (2007) and Miądlikowska et al. (2007).

Exemplar genera: *Candelaria* A. Massal. 1853, *Candelariella* Müll. Arg. 1894.

Order: **Umbilicariales** Lumbsch, Hestmark & Lutzoni, **ord. nov.**
MycoBank no.: MB 501293

Ascomycetes lichenisati algas virides thallo continentis. Ascumata sessilia, raro immersa usque ad paucum elevata, plerumque atra, irregularia, disciformia. Excipulum pigmentatum, annulatum. Hymenium amyloideum. Paraphyses simplices vel paulum ramosae, septatae. Asci unitunicati, tholo inconspicue amyloideo, 1–8-sporei. Ascosporeae hyalinae vel brunneae, non-septatae usque ad muriformes.

Typus: *Umbilicaria* Hoffm. 1789.

Lichen-forming ascomycetes with chlorococcoid photobiont. Ascumata sessile, or rarely immersed or stalked, mostly black, irregular, disciform. True exciple pigmented, annulate. Hymenium amyloid. Paraphyses simple or slightly branched, septate, apically thickened. Asci unitunicate, with slightly amyloid tholi, 1–8-spored. Ascospores hyaline or brown, non-septate to muriform.

Exemplar genera: *Lasallia* Mérat 1821, *Umbilicaria* Hoffm. 1789.

Class: **Leotiomyces** O. E. Erikss. & Winka, *Myconet* 1: 7 (1997).
Excluding *Geoglossaceae* (Wang et al. 2006).

Order: **Cyttariales** Luttr. ex Gamundí, *Darwiniana* 16: 502 (1971).

Synonym: *Cyttariales* Luttr., *Univ. Miss. Stud.* 24(2): 109 (1951), *nomen invalidum*.

Exemplar genus: *Cyttaria* Berk. 1842.

Order: **Erysiphales** H. Gwynne-Vaughan, *Fungi, Ascom., Ustilag.*, *Ured.*: 78 (1922).

Exemplar genera: *Erysiphe* R. Hedw. ex DC. 1805, *Blumeria* Golovin ex Speer 1975, *Uncinula* Lév. 1851.

Order: **Helotiales** Nannf., *Nova Acta R. Soc. Scient. Upsal.*, ser. 4 8(2): 68 (1932).

Based on current character and taxon sampling (Wang et al. 2006, 2007; Spatafora et al. 2007), the monophyly of *Helotiales* s. lat. is not well supported. There exists a minimum of five helotialean lineages that are intermixed with other leotiomyces taxa (e.g. *Cyttariales*, *Erysiphales*) resulting in a paraphyletic *Helotiales* s. lat. The interrelationships of these taxa are poorly resolved, however, thus preventing the synthesis of an accurate phylogenetic classification at this time. *Leotiomyces* represents one of the more undersampled higher taxa among the *Ascomycota*, and it is likely that future sampling will result in a phylogenetic classification of a more restricted *Helotiales* and the recognition of additional orders based on current helotialean families (e.g. *Leotiaceae* or *Helotiaceae*, *Sclerotiniaceae*).

Exemplar genera: *Mitruia* Fr. 1821, *Hymenoscyphus* Gray 1821, *Ascocoryne* J.W. Groves & D.E. Wilson 1967.

Order: **Rhytismatales** M. E. Barr ex Minter, in Hawksworth & Eriksson, *Syst. Ascom.* 5: 182 (1986).

Synonym: *Rhytismatales* M. E. Barr, *Mem. N. Y. Bot. Gdn* 28: 6 (1976), *nomen invalidum*.

Exemplar genera: *Rhytisma* Fr. 1818, *Lophodermium* Chevall. 1826, *Cudonia* Fr. 1849.

Order: **Thelebolales** P. F. Cannon, in Kirk et al., *Ainsworth & Bisby's Dict. Fungi* (9th edn): xi (2001).

Exemplar genera: *Thelebolus* Tode 1790, *Coprotus* Korf ex Korf & Kimbr. 1967, *Ascozonus* (Renny) E.C. Hansen 1876.

Class: **Lichinomycetes** Reeb, Lutzoni & Cl. Roux., *Mol. Phylogen. Evol.* 32: 1055 (2004).

Order: **Lichinales** Henssen & Büdel, in Hawksworth & Eriksson, *Syst. Ascom.* 5: 138 (1986).

Exemplar genera: *Heppia* Nägeli ex A. Massal. 1854, *Lichina* C. Agardh 1817, *Peltula* Nyl. 1853.

Class: **Orbiliomycetes** O. E. Erikss. & Baral, in Eriksson et al., *Myconet* 9: 96 (2003).

Order: **Orbiliales** Baral, O. E. Erikss., G. Marson & E. Weber, in Eriksson et al., *Myconet* 9: 96 (2003).

Exemplar genera: *Orbilium* Fr. 1849, *Hyalorbilia* Baral & G. Marson 2000.

Class: **Pezizomycetes** O. E. Erikss. & Winka, *Myconet* 1: 8 (1997).

Order: **Pezizales** J. Schröt., in Engler & Prantl (eds), *Nat. Pflanzenfam.* 1: 173 (1894), as 'Pezizineae'.

Exemplar genera: *Peziza* Fr. 1822, *Glaziella* Berk. 1880, *Morchella* Dill. ex Pers. 1794, *Pyronema* Carus 1835, *Tuber* F.H. Wigg. 1780.

Glaziella has been described several times, *inter alia* as a zygomycete. Gibson *et al.* (1986) demonstrated it was an ascomycete and proposed a new family and order close to Pezizales, but small subunit rRNA gene sequences show that it should be included in Pezizales (Landvik & Eriksson 1994).

Class: **Sordariomycetes** O. E. Erikss. & Winka, *Myconet* 1: 10 (1997).

Subclass: **Hypocreomycetidae** O. E. Erikss. & Winka, *Myconet* 1: 6 (1997).

Order: **Coronophorales** Nannf., *Nova Acta R. Soc. Scient. Upsal.*, ser. 4 8: 54 (1932).

Exemplar genera: *Nitschkia* G.H. Otth ex P. Karst. 1873, *Scortechinia* Sacc. 1885, *Bertia* De Not. 1844, *Chaetosphaerella* E. Müll. & C. Booth 1972.

Order: **Hypocreales** Lindau, in Engler & Prantl (eds), *Nat. Pflanzenfam.* 1: 343 (1897).

Exemplar genera: *Hypocrea* Fr. 1825, *Nectria* (Fr.) Fr. 1849, *Corcydiceps* (Fr.) Link 1833, *Claviceps* Tul. 1853, *Niesslia* Auersw. 1869.

Order: **Melanosporales** N. Zhang & M. Blackw., **ord. nov.**
MycoBank no.: MB 501294

Ascomata perithecialia vel nonnumquam ostiolo carentia; peridium ascomatis e basi glomeris ascogonialis oriundum, translucidum; centrum pseudoparenchymaticum, paraphysibus absentibus; asci unitunicati, evanescentes; ascosporae fuscae, poro germinationis utrinque praeditae; anamorphae hyphomycetales. Fungi saepe mycoparasitici.

Typus: *Melanospora* Corda 1837.

Ascoma perithecial or secondarily cleistothecial, peridium derived from base of an ascogonial coil, translucent; centrum pseudoparenchymatous, paraphyses absent in development; asci unitunicate, evanescent; ascospores dark, with germ pores at both ends; anamorphs hyphomycetous; often mycoparasitic.

Exemplar genus: *Melanospora* Corda 1837.

Order: **Microascales** Luttr. ex Benny & Kimbr., *Mycotaxon* 12: 40 (1980).

Synonym: *Microascales* Luttr., *Univ. Miss. Stud.* 24(2): 108 (1951), *nomen invalidum*.

The group as recognized here includes members of the *Halosphaeriales*. In Zhang *et al.* (2007) and Tang *et al.* (2007), the *Halosphaeriales* were maintained separate from the *Microascales*.

Exemplar genera: *Microascus* Zukal 1885, *Petriella* Curzi 1930, *Halosphaeria* Linder 1944, *Lignicola* Höhnk 1955, *Nimbospora* J. Koch 1982.

Subclass: **Sordariomycetidae** O. E. Erikss. & Winka, *Myconet* 1: 10 (1997).

Order: **Boliniales** P. F. Cannon, in Kirk *et al.*, *Ainsworth & Bisby's Dict. Fungi* (9th edn): x (2001).

Exemplar genera: *Camarops* P. Karst. 1873, *Apiocamarops* Samuels & J. D. Rogers 1987.

Order: **Calosphaeriales** M. E. Barr, *Mycologia* 75: 11 (1983).

This order has not been placed in a subclass but the work of Réblová *et al.* (2004) shows that it may be related to the *Diaporthales*. Members of this group were not included in Zhang *et al.* (2007) or Tang *et al.* (2007).

Exemplar genera: *Calosphaeria* Tul. & C. Tul. 1863, *Togniniella* Réblová, L. Mostert, W. Gams & Crous 2004, *Pleurostoma* Tul. & C. Tul. 1863.

Order: **Chaetosphaeriales** Huhndorf, A. N. Mill. & F. A. Fernández, *Mycologia* 96: 378 (2004).

Exemplar genera: *Chaetosphaeria* Tul. & C. Tul. 1863, *Melanochaeta* E. Müll., Harr & Sulmont 1969, *Zignoëlla* Sacc. 1878, *Striatosphaeria* Samuels & E. Müll. 1979.

Order: **Coniochaetales** Huhndorf, A. N. Mill. & F. A. Fernández, *Mycologia* 96: 378 (2004a).

Exemplar genera: *Coniochaeta* (Sacc.) Cooke 1887, *Coniochaetidium* Malloch & Cain 1971.

Order: **Diaporthales** Nannf., *Nova Acta R. Soc. Scient. upsal.*, ser. 4 8: 53 (1932).

Exemplar genera: *Diaporthe* Nitschke 1870, *Gnomonia* Ces. & De Not. 1863, *Cryphonectria* (Sacc.) Sacc. & D. Sacc. 1905, *Valsa* Fr. 1849.

Order: **Ophiostomatales** Benny & Kimbr., *Mycotaxon* 12: 48 (1980).

Exemplar genera: *Ophiostoma* Syd. & P. Syd. 1919, *Fragosphaeria* Shear 1923.

Order: **Sordariales** Chadeff. ex D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 182 (1986).

Synonym: *Sordariales* Chadeff., in Chadeffaud & Emberger, *Traité Bot.* 1: 594 (1960), *nomen invalidum*.

Exemplar genera: *Sordaria* Ces. & De Not. 1863, *Podospora* Ces. 1856, *Neurospora* Shear & B.O. Dodge 1927, *Lasiosphaeria* Ces. & De Not. 1863, *Chaetomium* Kunze 1817.

Subclass: **Xylariomycetidae** O. E. Erikss. & Winka, *Myconet* 1: 12 (1997).

Order: **Xylariales** Nannf., *Nova Acta R. Soc. Scient. Upsal.*, ser. 4, 8: 66 (1932).

Exemplar genera: *Xylaria* Hill ex Schrank 1789, *Hypoxylon* Bull. 1791, *Anthostomella* Sacc. 1875, *Diatrype* Fr. 1849, *Graphostroma* Piroz. 1974.

Sordariomycetes incertae sedis (not placed in any subclass)

Order: **Lulworthiales** Kohlm., Spatafora & Volkm-Kohlm., *Mycologia* 92: 456 (2000).

This order includes members formerly placed in the Spathulosporales.

Exemplar genera: *Lulworthia* G. K. Sutherl. 1916, *Lindra* I.M. Wilson 1956.

Order: **Meliolales** Gäum. ex D. Hawksw. & O. E. Erikss., *Syst. Ascom.* 5: 180 (1986).

Synonym: *Meliolales* Gäum., *Pilze* (2nd edn): 158 (1964), *nomen invalidum*.

Exemplar genus: *Meliola* Fr. 1825.

Order: **Phyllachorales** M. E. Barr, *Mycologia* 75: 10 (1983).

Exemplar genus: *Phyllachora* Nitschke ex Fuckel 1870.

Order: **Trichosphaeriales** M. E. Barr, *Mycologia* 75: 11 (1983).

Exemplar genus: *Trichosphaeria* Fuckel 1870.

Pezizomycotina incertae sedis (not placed in any class)

Order: **Lahmiales** O. E. Erikss., *Mycotaxon* 27: 357 (1986).

Exemplar genus: *Lahmia* Körb. 1861.

Order: **Medeolariales** Korf, in Eriksson *Mycotaxon* 15: 232 (1982).

Exemplar genus: *Medeolaria* Thaxt. 1922.

Order: **Triblidiales** O. E. Erikss., *Syst. Ascom.* 11: 9 (1992).

Exemplar genera: *Huangshania* O. E. Erikss. 1992, *Pseudoglyphis* Nyl. 1855, *Triblidium* Rebert. 1804.

Phylum: **Basidiomycota** R. T. Moore, *Bot. Mar.* 23: 371 (1980).

Synonyms: *Basidiomycota* Bold, *Morph. Pl.*: 7, 198 (1958), *nomen invalidum*;

Basidiomycetes Whittaker (1959: 220), *nomen invalidum*. (Table 3, Fig 3) As in the case of *Fungi*, Moore (1980) validated a name that had already been used by Bold (1957), but he did not cite Bold's work.

Subphylum: **Pucciniomycotina** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 45 (2006).

Equivalent to *Urediniomycetes* (Kirk et al. 2001; Swann & Taylor 1995; Swann et al. 2001). The classification of *Pucciniomycotina* employed here parallels that of Bauer et al. (2006) and Aime et al. (2007).

Class: **Pucciniomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 48 (2006).

Equivalent to *Urediniomycetidae* (Swann et al. 2001).

Order: **Septobasidiales** Couch ex Donk, *Persoonia* 3: 243 (1964).

Synonym: *Septobasidiales* Couch, *Gen. Septobasidium*: 65 (1938), *nomen invalidum*.

Exemplar genera: *Septobasidium* Pat. 1892, *Auriculosocypha* D. A. Reid & Manim. 1985.

Order: **Pachnocybales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 48 (2006).

Exemplar genus: *Pachnocybe* Berk. 1836.

Order: **Helicobasidiales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 48 (2006).

Exemplar genera: *Helicobasidium* Pat. 1885, *Tuberculina* Tode ex Sacc. 1880.

Order: **Platyglloeales** R. T. Moore, *Mycotaxon* 39: 247 (1990).

Equivalent to *Platyglloeales* s. str. (Swann et al. 2001).

Exemplar genera: *Platyglloea* J. Schröt. 1887 s. str., *Eocronartium* G.F. Atk. 1902.

Order: **Pucciniales** Clem. & Shear, *Gen. Fungi* (2nd edn): 147 (1931).

Equivalent to *Uredinales*.

Exemplar genera: *Puccinia* Pers. 1801, *Uromyces* (Link) Unger 1832.

Class: **Cystobasidiomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 46 (2006).

Equivalent to the *Erythrobasidium-Naohidea-Sakaguchia* clade (Swann et al. 2001) and *Cystobasidiaceae* lineage (Weiß et al. 2004a). Genera of *Cystobasidiomycetes* that are not placed in any order include *Sakaguchia* Y. Yamada, K. Maeda & Mikata 1994, and *Cyrenella* Goch. 1981 (Aime et al. 2007; Bauer et al. 2006).

Order: **Cystobasidiales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 46 (2006).

Exemplar genera: *Cystobasidium* (Lagerh.) Neuhoff 1924, *Occultifur* Oberw. 1990, *Rhodotorula* F.C. Harrison 1927 *pro parte*.

Order: **Erythrobasidiales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 46 (2006).

Exemplar genera: *Erythrobasidium* Hamam. Sugiyama & Komag. 1988, *Rhodotorula* F. C. Harrison 1927 *pro parte*, *Sporobolomyces* Kluyver & C. B. Niel 1924 *pro parte*, *Bannoa* Hamam. 2002.

Order: **Naohideales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 46 (2006).

Exemplar genus: *Naohidea* Oberw. 1990.

Class: **Agaricostilbomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 45 (2006).

Equivalent to *Agaricostilbomycetidae* (Swann et al. 2001; Weiß et al. 2004a).

Order: **Agaricostilbales** Oberw. & R. Bauer, *Sydowia* 41: 240 (1989).

Exemplar genera: *Agaricostilbum* J. E. Wright 1970 (emend. Wright, Bandoni & Oberw. 1981), *Chionosphaera* D. E. Cox 1976, *Kondoa* Y. Yamada, Nakagawa & I. Banno 1989 (emend. Fonseca, Sampaio, Inácio & Fell 2000).

Order: **Spiculogloeales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 45 (2006).

Equivalent to *Mycogloea* group (Weiß et al. 2004a).

Exemplar genera: *Mycogloea* L. S. Olive 1950, *Spiculogloea* P. Roberts 1996, *Sporobolomyces* Kluyver & C. B. Niel 1924 *pro parte*.

Table 3 – Support for major groups of Fungi in selected phylogenetic studies: Basidiomycota

Rank	Taxon	Reference	Data	OTUs	Support
Phylum	BASIDIOMYCOTA	James et al. (2006)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	50	BPP = 1 MLBS = 80
Subphylum	Pucciniomycotina	Matheny et al. (2007a, fig. 4)	SSU, LSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	17	BPP = 1 MPBS = 100
		Aime et al. (2007, fig. 2)	LSU, SSU	109	BPP = 1 MPBS = 100
Class	Pucciniomycetes	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	7	BPP > 0.95 MPBS > 70
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	24	BPP = 0.97 MPBS ≥ 70
		Aime et al. (2007, fig. 2)	LSU, SSU	19	BPP = 1 MPBS = 100
		Aime et al. (2007, fig. 3)	LSU, SSU	41	MPBS = 86
Order	Septobasidiales	Arun Kumar et al. (2007, fig. 7)	LSU, SSU	4	BPP = 1 MPBS = 100
Order	Pachnocybales	Bauer et al. (2006, fig. 1)	LSU	1	NA
		Berres et al. (1995, fig. 4)	LSU	1	NA
Order	Helicobasidiales	Aime et al. (2007, fig. 2)	LSU, SSU	2	BPP = 1 MPBS = 96 NJBS = 98
		Aime et al. (2007, fig. 3)	LSU, SSU	10	MPBS = 87
Order	Platyglloeales	Aime et al. (2007, fig. 2)	LSU, SSU	4	BPP = 1 MPBS = 100 NJBS = 100
		Aime et al. (2007, fig. 3)	LSU, SSU	8	MPBS = 99
		Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	2	BPP > 0.95 MPBS > 70
Order	Pucciniales	Aime et al. (2007, fig. 2)	LSU, SSU	12	BPP = 1 MPBS = 100 NJBS = 100
		Aime (2006)	LSU	46	MPBS = 99
		Wingfield et al. (2004)	SSU	72	MPBS < 50
		Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	5	BPP > 0.95 MPBS > 70
Class	Cystobasidiomycetes	Aime et al. (2007, fig. 2)	LSU, SSU	27	BPP = 1 MPBS = 100 NJBS = 96
		Sampaio (2004, fig. 1)	LSU	11	BPP = 0.92
		Sampaio (2004, fig. 2)	LSU	26	BPP = 0.98
Order	Cystobasidiales	Nagahama et al. (2006, fig. 2)	LSU, SSU, 5.8S, <i>tef1</i>	9	MLBS = 100
		Aime et al. (2007, fig. 2)	LSU, SSU	12	BPP = 1 MPBS = 100 NJBS = 100
		Sampaio (2004, fig. 2)	LSU	8	BPP = 1
Order	Erythrobasidiales	Nagahama et al. (2006, fig. 2)	LSU, SSU, 5.8S, <i>tef1</i>	21	MLBS = 72
		Aime et al. (2007, fig. 2)	LSU, SSU	14	BPP = 1 MPBS = 83 NJBS = 91
		Sampaio (2004, fig. 2)	LSU	18	BPP = 1
Order	Naohideales	Aime et al. (2007, fig. 3)	LSU, SSU	2	MPBS = 98
		Weiß et al. (2004)	LSU	3	BPP = 0.94 NJBS < 50
Class	Agaricostilbomycetes	Aime et al. (2007, fig. 2)	LSU, SSU	25	BPP = 1 MPBS < 70 NJBS < 70
		Bauer et al. (2006, fig. 2)	LSU, SSU	4	NJBS = 89
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	8	BPP = 1 MPBS > 70
Order	Agaricostilbales	Aime et al. (2007, fig. 2)	LSU, SSU	22	BPP = 1 MPBS = 100 NJBS = 100
		Aime et al. (2007, fig. 2)	LSU, SSU	34	MPBS = 98
		Sampaio (2004, fig. 1)	LSU	7	BPP = 1
		Sampaio (2004, fig. 2)	LSU	23	BPP = 1

(continued on next page)

Table 3 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
Order	Spiculogloales	Fell et al. (2001)	LSU	24	MPBS = 64
		Aime et al. (2007, fig. 2)	LSU, SSU	3	BPP = 1 MPBS = 100 NJBS = 100
Class	Microbotryomycetes	Aime et al. (2007, fig. 3)	LSU, SSU	7	MPBS = 74
		Bauer et al. (2006, fig. 2)	LSU, SSU	2	NJBS = 90
		Aime et al. (2007, fig. 2)	LSU, SSU	31	BPP = 1 MPBS = 100 NJBS = 100
		Aime et al. (2007, fig. 3)	LSU, SSU	60	MPBS = 74
Order	Heterogastridiales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	6	BPP > 0.95 MPBS > 70
		Sampaio (2004, fig. 2)	LSU	49*	BPP = 0.87
Order	Heterogastridiales	Fell et al. (2001)	LSU	78	MPBS = 75
Order	Microbotryales	Bauer et al. (2006, fig. 2)	LSU, SSU	1	NA
Order	Microbotryales	Aime et al. (2007, fig. 2)	LSU, SSU	4	BPP = 1 MPBS = 99 NJBS = 94
Order	Leucosporidiales	Aime et al. (2007, fig. 3)	LSU, SSU	12	MPBS = 82
		Aime et al. (2007, fig. 2)	LSU, SSU	3	BPP = 0.98 MPBS = 85 NJBS = 100
Order	Sporidiobolales	Aime et al. (2007, fig. 3)	LSU, SSU	9	MPBS = 67
		Aime et al. (2007, fig. 2)	LSU, SSU	13	BPP = 1 MPBS = 74 NJBS = 68
Class/Order	Atractiellomycetes, Atractiellales	Aime et al. (2007, fig. 3)	LSU, SSU	17	MPBS = 69
		Sampaio (2004, fig. 2)	LSU	20	BPP = 0.98
		Aime et al. (2007, fig. 2)	LSU, SSU	4	BPP = 1 MPBS = 80 NJBS = 96
Class/Order	Classiculomycetes, Classiculales	Aime et al. (2007, fig. 3)	LSU, SSU	8	MPBS = 68
		Bauer et al. (2006, fig. 2)	LSU, SSU	7	NJBS = 68
		Aime et al. (2007, fig. 2)	LSU, SSU	2	BPP = 1 MPBS = 100 NJBS = 100
Class/Order	Mixiomycetes, Mixiales	Weiß et al. (2004, figs. 1–2)	LSU	2	BPP = 1 NJBS = 99
		Aime et al. (2007, fig. 2)	LSU, SSU	1	NA
Class/Order	Cryptomycocolacomycetes, Cryptomycocolacales	Bauer et al. (2006, fig. 2)	LSU, SSU	1	NA
		Aime et al. (2007, fig. 3)	LSU, SSU	1	NA
Subphylum	Ustilaginomycotina	Bauer et al. (2006, fig. 1)	LSU	2	NJBS = 100
		Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	24	BPP = 1 MPBS = 100
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	59	BPP = 1 MPBS > 70
Class	Ustilaginomycetes	Bauer et al. (2006, fig. 2)	LSU, SSU	21	NJBS = 100
		Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	12	BPP > 0.95 MPBS > 70
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	25	BPP = 1 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	53	BPP = 1 MPBS = 83 NJBS = 77
Order	Urocystales	Bauer et al. (2001, figs. 33–34)	LSU	36	MPBS = 79 NJBS = 93
		Fell et al. (2001, fig. 24)	LSU	27	NJBS = 86
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	5	BPP = 1 MPBS = 66 NJBS = 96
		Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	1	NA
		Bauer et al. (2001, figs. 33–34)	LSU	9	MPBS = 95 ³ NJBS = 96 ³

Table 3 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
Order	Ustilaginales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	10	BPP > 0.95 MPBS > 70
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	23	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	46	BPP = 1 MPBS < 60 NJBS < 60
Class	Exobasidiomycetes	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	12	BPP > 0.95 MPBS < 50
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	35	BPP < 0.60 MPBS < 60 NJBS < 60
		Bauer et al. (2001, figs. 33–34)	LSU	36	MPBS = 85 NJBS = 56
Order	Doassansiales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	1	NA
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	4	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	4	BPP = 1 MPBS = 84 NJBS = 77
		Bauer et al. (2001, figs. 33–34)	LSU	5	MPBS = 96 NJBS = 97
Order	Entylomatales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	4	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	3	BPP = 1 MPBS < 60 NJBS < 60
		Bauer et al. (2001, figs. 33–34)	LSU	9	MPBS = 72 NJBS = 91
Order	Exobasidiales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	2	BPP > 0.95 MPBS > 70
		Matheny et al. (2007a, fig. 5)	LSU, SSU	6	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	8	BPP = 1 MPBS < 60 NJBS = 61
Order	Georgefischeriales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	2	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	5	BPP < 0.60 MPBS < 60 NJBS < 60
		Bauer et al. (2001, figs. 33–34)	LSU	9	MPBS = 86 NJBS = 65
Order	Microstromatales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	1	NA
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	3	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	5	BPP = 1 MPBS = 63 NJBS = 67
Order	Tilletiales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	2	BPP > 0.95 MPBS > 70
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	7	BPP > 0.95 MPBS > 70
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	5	BPP = 1 MPBS = 76 NJBS = 64
Order	Ustilaginomycotina incertae sedis (not placed in any class) Malasseziales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	1	NA
		Begerow et al. (2007, fig. 1)	LSU, ITS, <i>atp6</i> , <i>βtub</i>	2	BPP = 1 MPBS = 100 NJBS = 100

(continued on next page)

Table 3 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
		Bauer et al. (2001, figs. 33–34)	LSU	4	MPBS = 100 NJBS = 100
Subphylum	Agaricomycotina	Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	125	BPP = 1 MPBS = 95
Class	Tremellomycetes	Matheny et al. (2007a, fig. 4)	SSU, LSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	5	BPP > 0.95 MPBS = 50–69
Order	Cystofilobasidiales	Fell et al. (2001, figs. 19, 22)	LSU	139	MPBS = 100
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	5	BPP = 1 MPBS ≥ 70
		Fell & Scorzetti (2004, fig. 1)	LSU	16	BPP = 1 MPBS = 83
Order	Filobasidiales	Fell et al. (2001, figs. 19, 22)	LSU	34	MPBS = 96
Order	Tremellales	Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	5	BPP ≥ 0.95 MPBS ≥ 70
Class/Order	Dacrymycetes, Dacrymycetales	Fell et al. (2001, figs. 19, 22)	LSU	89	MPBS = 56
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	4	BPP = 1 MPBS = 100
Class	Agaricomycetes	Weiß & Oberwinkler (2001, fig. 6)	LSU	9	NJBS = 99
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	119	BPP = 1 MPBS = 95
		James et al. (2006)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	37	BPP = 1 MLBS = 92
Subclass	Agaricomycetidae	Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	63	BPP = 1 MPBS = 96
		Binder & Hibbett (2007, fig. 2)	LSU, SSU, 5.8S, mt-LSU, <i>atp6</i>	47	BPP > 0.98 MLBS = 88
Order	Agaricales	Binder et al. (2005, fig. 1)	LSU, SSU, mt-LSU, mt-SSU	46	MPBS = 62
		Matheny et al. (2006, fig. 2)	LSU, SSU, 5.8S	230	BPP = 0.84
		Matheny et al. (2006, fig. 3)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i>	238	BPP = 1 MPBS = 43
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	41	BPP = 1 MPBS = 76
Order	Atheliales	Moncalvo et al. (2002, fig. 2)	LSU	786	MPBS < 50
		Larsson et al. (2004, fig. 1)	LSU	8	MPBS = 97
Order	Boletales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	3	MPBS = 75
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	11	BPP = 1 MPBS = 100
		Binder & Hibbett (2007, fig. 2)	LSU, SSU, 5.8S, mt-LSU, <i>atp6</i>	42	BPP > 0.98 MLBS = 99
Subclass	Phallomycetidae	Binder & Hibbett (2007, fig. 3)	LSU	301	BPP > 0.98
		Hosaka et al. (2007, fig. 2)	LSU, mt-SSU, <i>atp6</i> , <i>rpb2</i> , <i>tef1</i>	222	BPP = 1 MPBS = 98
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	3	BPP = 1 MPBS = 100
Order	Geastrales	Hosaka et al. (2007, fig. 2)	LSU, mt-SSU, <i>atp6</i> , <i>rpb2</i> , <i>tef1</i>	21	BPP = 1 MPBS = 59
Order	Gomphales	Hosaka et al. (2007, fig. 2)	LSU, mt-SSU, <i>atp6</i> , <i>rpb2</i> , <i>tef1</i>	61	BPP = 1 MPBS = 63
Order	Hysterangiales	Hosaka et al. (2007, fig. 2)	LSU, mt-SSU, <i>atp6</i> , <i>rpb2</i> , <i>tef1</i>	99	BPP = 1 MPBS = 98
Order	Phallales	Hosaka et al. (2007, fig. 2)	LSU, mt-SSU, <i>atp6</i> , <i>rpb2</i> , <i>tef1</i>	41	BPP = 1 MPBS = 84
	Agaricomycetes incertae sedis (not placed in any subclass):				
Order	Auriculariales	Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	3	BPP = 1 MPBS = 100
Order	Cantharellales	Weiß & Oberwinkler (2001, fig. 6)	LSU	43	NJBS < 60
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	11	BPP = 1 MPBS = 69
		Moncalvo et al. (2007, fig. 1)	LSU, SSU, mtSSU, <i>rpb2</i>	29	BPP < 0.50 MPBS < 50
Order	Corticiales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	31	MPBS < 50
		Larsson et al. (2004, fig. 1)	LSU	7	MPBS = 96

Table 3 (continued)

Rank	Taxon	Reference	Data	OTUs	Support
Order	Gloeophyllales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	8	MPBS = 81
		Thorn et al. (2000, fig. 5)	LSU	5	MPBS = 71
	Hymenochaetales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	6	MPBS = 54
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	7	BPP = 1 MPBS = 63
Order	Polyporales	Larsson et al. (2007, fig. 3)	LSU, 5.8S	174	BPP = 1
		Wagner & Fischer (2002, fig. 2)	LSU	104	NJBS = 100
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	16	BPP = 1 MPBS = 85
Order	Russulales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	122	MPBS < 50
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	8	BPP = 1 MPBS = 99
Order	Sebacinales	Larsson & Larsson (2003, fig. 1)	LSU, 5.8S	127	MPBS = 96
		Miller et al. (2007, fig. 2)	LSU, ITS	143	MPBS = 100
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	2	BPP = 1 MPBS = 100
Order	Thelephorales	Weiß & Oberwinkler (2001, fig. 6)	LSU	9	NJBS = 99
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	2	BPP = 1 MPBS = 100
Order	Trechisporales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	13	MPBS = 97
		Larsson et al. (2004, fig. 1)	LSU, 5.8S	11	MPBS = 86
		Matheny et al. (2007b, fig. 6)	LSU, SSU, 5.8S, <i>rpb2</i> , <i>tef1</i>	2	BPP = 1 MPBS = 100
Class/Order	Basidiomycota incertae sedis (not placed in any subphylum): Wallemiomycetes, Wallemiales	Binder et al. (2005, fig. 4)	LSU, SSU, mt-LSU, mt-SSU	20	MPBS = 69
		Larsson et al. (2004, fig. 1)	LSU, 5.8S	12	MPBS = 99
Class/Order	Entorrhizomycetes, Entorrhizales	Matheny et al. (2007a, fig. 4)	LSU, SSU, 5.8S, <i>rpb1</i> , <i>rpb2</i> , <i>tef1</i>	3	BPP > 0.95 MPBS > 70
		Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	3	BPP = 1 MPBS > 70
Class/Order	Entorrhizomycetes, Entorrhizales	Matheny et al. (2007a, fig. 5)	LSU, SSU, 5.8S	3	BPP = 1 MPBS > 70
		Bauer et al. (2001, figs. 33–34)	LSU	2	MPBS = 100 NJBS = 100

See Table 1 for explanation.

Class: **Microbotryomycetes** R. Bauer, Begerow, J. P. Sampaio, M. Weiß & Oberw., *Mycol. Progr.* 5: 47 (2006).

Equivalent to *Microbotryomycetidae* (Swann et al. 2001; Weiß et al. 2004a). The backbone of the *Microbotryomycetes* remains poorly resolved, and several genera of *Microbotryomycetes* are not placed in any order, including *Colacogloea* Oberw. & R. Bauer 1991, *Atractocolax* R. Kirschner, R. Bauer & Oberw. 1999, *Krieglsteinera* Pouzar 1987, *Camptobasidium* Marvanová & Suberkr. 1990, *Kriegeria* Bres. 1891 and certain species of the polyphyletic genera *Sporobolomyces* Kluyver & C. B. Niel 1924 *pro parte*, *Rhodotorula* F. C. Harrison 1927 *pro parte*, and *Leucosporidium* Fell, Statzell, I. L. Hunter & Phaff 1970, and others (Aime et al. 2007; Bauer et al. 2006; Sampaio et al. 2003; Weiß et al. 2004a).

Order: **Heterogastridiales** Oberw., R. Bauer & Bandoni R. J., *Mycologia* 82: 57 (1990).

Exemplar genus: *Heterogastridium* Oberw. & R. Bauer 1990.

Bauer et al. (2006) placed *Colacogloea*, *Atractocolax* and *Krieglsteinera* in the *Heterogastridiales*. However, analyses of Bauer

et al. (2006) and Aime et al. (2007) suggest that *Heterogastridium* and *Colacogloea* do not form a clade, while *Atractocolax* and *Krieglsteinera* have yet to be sampled in molecular phylogenetic studies.

Order: **Microbotryales** R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* 75: 1309 (1997).

Exemplar genera: *Microbotryum* Lév. 1847, *Ustilentyloma* Savile 1964.

Order: **Leucosporidiales** J. P. Sampaio, M. Weiß & R. Bauer, in Sampaio et al., *Mycol. Progr.* 2: 61 (2003).

Exemplar genera: *Leucosporidiella* J. P. Sampaio 2003, *Leucosporidium* Fell, Statzell, I. L. Hunter & Phaff 1970, *Mastigobasidium* Golubev 1999.

Order: **Sporidiobolales** J. P. Sampaio, M. Weiß & R. Bauer, in Sampaio et al., *Mycol. Progr.* 2: 66 (2003).

Exemplar genera: *Sporidiobolus* Nyland 1949, *Sporobolomyces* Kluyver & C. B. Niel 1924, *Rhodosporeidium* I. Banno 1967, *Rhodotorula* F. C. Harrison 1927 *pro parte*.




Fig 3 – Phylogeny and classification of Fungi. Basidiomycota. See Table 3 for support values for clades. Dashed lines indicate taxa that are of uncertain placement.

Class: **Atractiellomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 45 (2006).

Order: **Atractiellales** Oberw. & Bandoni, *Can. J. Bot.* 60: 1740 (1982).

Emend. Oberw. & Bauer, *Sydowia* 41: 239 (1989).

Exemplar genera: *Atractiella* Sacc. 1886, *Saccoblastia* A. Møller 1895, *Helicogloea* Pat. 1892, *Phleogena* Link 1833.

Class: **Classiculomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 46 (2006).

Order: **Classicuales** R. Bauer, Begerow, Oberw. & Marvanová, *Mycologia* 95: 763 (2003).

Exemplar genera: *Classicula* R. Bauer, Begerow, Oberw. & Marvanová 2003, *Jaculispora* H. J. Huds. & Ingold 1960.

Class: **Mixiomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 47 (2006).

Order: **Mixiales** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 47 (2006).

Exemplar genus: *Mixia* C. L. Kramer 1959 [‘1958’].

Class: **Cryptomycocolacomycetes** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 46 (2006).

Order: **Cryptomycocolacales** Oberw. & R. Bauer, *Mycologia* 82: 672 (1990).

Exemplar genera: *Cryptomycocolax* Oberw. & R. Bauer 1990, *Colacosiphon* R. Kirschner, R. Bauer & Oberw. 2001.

Subphylum: **Ustilaginomycotina** R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* 5: 45 (2006).

Equivalent to *Ustilaginomycetes* (Bauer et al. 1997, 2001; Swann & Taylor 1995).

The classification of *Ustilaginomycotina* employed here largely parallels that of Begerow et al. (2007), with the primary differences being that here the *Entorrhizomycetes* are classified as *incertae sedis* among *Basidiomycota* (rather than being a class within *Ustilaginomycotina*).

Class: **Ustilaginomycetes** R. Bauer, Oberw. & Vánky, *Can. J. Bot.* 75: 1311 (1997).

Emend. Begerow, Stoll & Bauer, *Mycologia* 98: 908 (2007) [‘2006’].

Equivalent to *Ustilaginomycetidae* Jülich as emended by Bauer & Oberwinkler (Bauer et al. 1997, 2001; Weiß et al. 2004a).

Order: **Urocystales** R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* 75: 1311 (1997).

Exemplar genera: *Urocystis* Rabenh. ex Fuckel 1870, *Ustacystis* Zundel 1945, *Doassansiopsis* (Setch.) Dietel 1897.

Melanotaenium de Bary 1874 has also been placed in this order (Bauer et al. 2001; Weiß et al. 2004a), but analyses of Begerow et al. (2007) and Matheny et al. (2007b) have supported its transfer to *Ustilaginales*.

Order: **Ustilaginales** G. Winter, *Rabenh. Krypt.-Fl.* 2nd ed. 1(1.1): 73 (1880), as ‘*Ustilagineae*’.

Emend. Bauer & Oberwinkler, in Bauer et al., *Can. J. Bot.* 75: 1311 (1997).

Exemplar genera: *Ustilago* (Pers.) Roussel 1806, *Cintractia* Cornu 1883.

Thecaphora Fingerh. 1836 has also been placed in this order (Bauer et al. 2001), but analyses of Begerow et al. (2007) and Matheny et al. (2007b) have suggested that it is not nested in *Ustilaginales*. *Thecaphora* may be the sister group of *Urocystales* (Matheny et al. 2007b).

Class: **Exobasidiomycetes** Begerow, Stoll & R. Bauer, *Mycologia* 98: 908 (2007) [‘2006’].

Equivalent to *Exobasidiomycetidae* Jülich 1981 *emend.* Bauer & Oberwinkler, except for exclusion of *Malasseziales* (Bauer et al. 1997, 2001; Weiß et al. 2004a).

Monophyly of the *Exobasidiomycetidae*, as delimited here, is supported with high Bayesian posterior probability in analyses of *rpb1*, *rpb2*, and *tef1*, and nuclear *lsu*, *ssu*, and 5.8S ribosomal genes (Matheny et al. 2007b), but it is weakly supported in analyses using *atp6*, β -tubulin, and nuclear *lsu* ribosomal RNA genes (Begerow et al. 2007). See comments regarding *Malasseziales*.

Order: **Doassansiales** R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* 75: 1312 (1997).

Exemplar genera: *Doassansia* Cornu 1883, *Rhamphospora* D. D. Cunn. 1888, *Nannfeldtiomyces* Vánky 1981.

Order: **Entylomatales** R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* **75**: 1311 (1997).

Exemplar genera: *Entyloma* de Bary 1874, *Tilletiopsis* Derx 1948.

Begerow et al. (2007) erected the monotypic order *Ceraceosorales* Begerow, Stoll & R. Bauer for *Ceraceosorus bombacis* (B. K. Bakshi) B. K. Bakshi 1976, which was weakly supported as the sister group of *Tilletiopsis albescens* Gokhale 1972. The *Ceraceosorus-T. albescens* clade was placed as the sister group of *Entylomatales*, again with weak support. *Ceraceosorales* is not included in the present classification, pending more robust resolution of the relationships among *Ceraceosorus*, *Tilletiopsis*, and *Entyloma*.

Order: **Exobasidiales** Henn., in Engler & Prantl (eds), *Nat. Pflanzenfam.* **1(1**)**: 103 (1897), as ‘*Exobasidiineae*’.

Emend. Bauer, Oberwinkler & Vánky, *Can. J. Bot.* **75**: 1312 (1997).

Exemplar genera: *Exobasidium* Woronin 1867, *Clinoconidium* Pat. 1898, *Dicellomyces* L. S. Olive 1945.

Order: **Georgefischeriales** R. Bauer, Begerow & Oberw., in Bauer et al., *Can. J. Bot.* **75**: 1311 (1997).

Exemplar genera: *Georgefischeria* Thirum. & Naras. *emend.* Gandhe 1980, *Phragmotenium* R. Bauer, Begerow, A. Nagler & Oberw. 2001, *Tilletiaria* Bandoni & B. N. Johri 1972, *Tilletiopsis* Derx 1948 *pro parte*.

Order: **Microstromatales** R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* **75**: 1311 (1997).

Exemplar genera: *Microstroma* Niessl 1861, *Sympodiomyces* Sugiy., Tokuyoko & Komag. 1991, *Volvocisporium* Begerow, R. Bauer & Oberw. 2001.

Order: **Tilletiales** Kreisel ex R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* **75**: 1311 (1997).

Exemplar genera: *Tilletia* Tul. & C. Tul. 1847, *Conidiosporomyces* Vánky 1992, *Erratomyces* M. Piepenbr. & R. Bauer 1997.

Ustilaginomycotina incertae sedis (not placed in any class):

Order: **Malasseziales** R. T. Moore, *Bot. Mar.* **23**: 371 (1980).

Emend. Begerow, Bauer & Boekhout, *Mycol. Res.* **104**: 59 (2000).

Exemplar genus: *Malassezia* Baill. 1889.

Analyses of the protein-coding genes *rpb1*, *rpb2*, and *tef1*, alone or in combination with nuclear LSU, SSU, and 5.8S ribosomal genes, suggest that *Malasseziales* are included in the *Ustilaginomycetes*, but analyses of nuclear ribosomal genes alone or in combination with *atp6* and β -tubulin suggest that *Malasseziales* is in the *Exobasidiomycetes* (Bauer et al. 2001; Begerow et al. 2007; Matheny et al. 2007b; Weiß et al. 2004a).

Subphylum: **Agaricomycotina** Dowell *Prosyllabus* LXXVII (2001).

Homonym: *Agaricomycotina* R. Bauer, Begerow, J. P. Samp., M. Weiß & Oberw., *Mycol. Progr.* **5**: 45 (2006). Equivalent to

Hymenomycetes (Swann & Taylor 1995) or *Basidiomycetes* (Kirk et al. 2001; Hibbett 2007).

Class: **Tremellomycetes** Dowell, *Prosyllabus*: LXXVII (2001).

Dimorphic fungi. Fruiting bodies gelatinous or absent, parentheses sacculate or absent, basidia septate or non-septate. The least inclusive clade containing *Tremellales*, *Filobasidiales* and *Cystofilobasidiales*.

Equivalent to *Tremellomycetidae* sensu Swann & Taylor (1995) and Weiß et al. (2004a). The name *Tremellomycetidae* was earlier published by Locquin (1984), but without a Latin diagnosis, and it is therefore invalid under the Code.

Order: **Cystofilobasidiales** Fell, Roelijmans & Boekhout, *Int. J. Syst. Bacteriol.* **49**: 911 (1999).

Exemplar genera: *Cystofilobasidium* Oberw. & Bandoni 1983, *Mrakia* Y. Yamada & Komag. 1987, *Itersonilia* Derx 1948.

Order: **Filobasidiales** Jülich, *Bibliotheca Mycol.* **85**: 324 (1981).

Exemplar genera: *Filobasidiella* Kwon-Chung 1976, *Cryptococcus* Vuill. 1901 (*pro parte*).

Order: **Tremellales** Fr., *Syst. Mycol.* **1**: 2 (1821), as ‘*Tremellinae*’.

As delimited here, the group includes *Trichosporonales* Boekhout & Fell 2001 (Fell et al. 2001) and *Christianseniales* F. Rath 1991 (Wells & Bandoni 2001). *Filobasidiales*, which Weiß et al. (2004a) included in *Tremellales* s. lat., has been resolved as the sister group of *Tremellales* (Fell et al. 2001; Matheny et al. 2007b; Swann & Taylor 1995).

Exemplar genera: *Tremella* Pers. 1794, *Trichosporon* Behrend 1890, *Christiansenia* Hauerslev 1969.

Class: **Dacrymycetes** Dowell, *Prosyllabus*: LXXVII (2001)

Fruiting bodies gelatinous, basidia furcate (rarely unisporous), parentheses imperforate.

Containing the single order *Dacrymycetales* (Wells & Bandoni 2001).

Order: **Dacrymycetales** Henn., in Engler & Prantl (eds), *Nat. Pflanzenfam.* **1(1**)**: 96 (1898), as ‘*Dacrymycetinae*’.

Exemplar genera: *Dacrymyces* Nees 1861, *Calocera* (Fr.) Fr. 1828, *Guepiniopsis* Pat. 1883.

Class: **Agaricomycetes** Dowell, *Prosyllabus*: LXXVII (2001)

Fruiting bodies hymenomycetous or gasteroid, basidia two- to eight-spored, parentheses perforate or imperforate. The least-inclusive clade containing *Auriculariales*, *Sebacinales*, *Cantharellales*, *Phallomycetidae* and *Agaricomycetidae*.

This group is approximately equivalent to *Homobasidiomycetes* sensu Hibbett & Thorn (2001) plus *Auriculariales* and *Sebacinales*.

Subclass: **Agaricomycetidae** Parmasto, *Windahlia* **16**: 16 (1986).
Synonym: *Basidiosporeae* Bessey, *Univ. Studies, Univ. Nebr.*

7: 306 (1907).

The least-inclusive clade containing *Agaricales*, *Boletales* and *Atheliales*.

The delimitation of *Agaricomycetidae* adopted here differs from that of Parmasto (1986), who described *Agaricomycetidae*

as a subclass of *Cantharellomycetes* Parm. 1986. For example, many of the resupinate forms in the *Agaricomycetidae* were placed by Parmasto in the *Corticomycetes* Parm. 1986. The name *Agaricomycetidae* was also published by Locquin (1984), but without a Latin diagnosis and it is therefore invalid under the Code.

Order: **Agaricales** Underw., *Moulds, Mildews Mushrooms*: 97 (1899).

Equivalent to euagarics clade (Hibbett & Thorn 2001).

Exemplar genera: *Agaricus* L. 1753, *Coprinus* Pers. 1797, *Pleurotus* (Fr.) P. Kumm. 1871.

Order: **Atheliales** Jülich, *Bibliotheca Mycol.* 85: 343 (1981).

Equivalent to athelioid clade (Binder et al. 2005; Larsson et al. 2004).

Exemplar genera: *Athelia* Pers. 1822, *Piloderma* Jülich 1969, *Tylospora* Donk 1960.

Order: **Boletales** E.-J. Gilbert, *Livres Mycol.* 3: 83 (1931).

Equivalent to bolete clade (Binder & Hibbett 2006; Hibbett & Thorn 2001).

Exemplar genera: *Boletus* Fr. 1821, *Scleroderma* Pers. 1801, *Coniophora* DC. 1815, *Rhizopogon* Fr. & Nordholm 1817.

Subclass: **Phallomycetidae** K. Hosaka, Castellano & Spatafora, *Mycologia* 98: 955 (2007) ['2006'].

Equivalent to *Phallales* sensu Kirk et al. (2001), and the gomphoid-phalloid clade (Hibbett & Thorn 2001; Hosaka et al. 2007).

Order: **Geastrales** K. Hosaka & Castellano, *Mycologia* 98: 957 (2007) ['2006'].

Exemplar genera: *Geastrum* Pers. 1794, *Radiigera* Zeller 1944, *Sphaerobolus* Tode 1790.

Order: **Gomphales** Jülich, *Bibliotheca Mycol.* 85: 348 (1981).

Exemplar genera: *Gomphus* (Fr.) Weinm. 1826, *Gautieria* Vittad. 1831, *Ramaria* Holmsk. 1790.

Order: **Hysterangiales** K. Hosaka & Castellano, *Mycologia* 98: 956 (2007) ['2006'].

Exemplar genera: *Hysterangium* Vittad. 1831, *Phallogaster* Morgan 1893, *Gallacea* Lloyd 1905, *Austrogautieria* E. L. Stewart & Trappe 1985.

Order: **Phallales** E. Fisch., in Engler & Prantl (eds), *Nat. Pflanzenfam.* 1(1**): 276 (1898).

Equivalent to *Phallomycetidae* Locq. (Locquin 1984), which was invalidly published, owing to the absence of a Latin diagnosis.

Exemplar genera: *Phallus* Junius ex L. 1753, *Clathrus* P. Micheli ex L. 1753, *Claustula* K. M. Curtis 1926.

Agaricomycetes incertae sedis (not placed in any subclass):

Order: **Auriculariales** J. Schröt., in Cohn (ed.), *Krypt.-Fl. Schlesien* 1: 382 (1889).

Exemplar genera: *Auricularia* Bull. ex Juss. 1789, *Exidia* Fr. 1822, *Bourdotia* (Bres.) Trotter 1913.

Order: **Cantharellales** Gäum., *Vergl. Morph. Pilze*: 495 (1926).

Equivalent to the cantharelloid clade (Hibbett & Thorn 2001; Moncalvo et al. 2007). The *Cantharellales* as delimited here includes *Tulasnella*, which is distinguished by unusual basidia with inflated sterigmata, and has been classified in a separate order, *Tulasnellales* Rea 1922 (e.g. Weiß et al. 2004a). Extreme evolutionary rate heterogeneity in the nuclear ribosomal RNA genes of *Tulasnella*, *Cantharellus* and *Craterellus* is a source of error in phylogenetics of *Cantharellales*. Analyses of Matheny et al. (2006) suggest that *Tulasnella* is nested within the *Cantharellales*, but it could also be the sister group to *Cantharellales* s.str. (Moncalvo et al. 2007). If so, then it may be appropriate to segregate *Tulasnellales* from *Cantharellales* s.str.

Exemplar genera: *Cantharellus* Fr. 1821, *Botryobasidium* Donk 1931, *Craterellus* Pers. 1825, *Tulasnella* J. Schröt. 1888.

Order: **Corticiales** K. H. Larss., *ord. nov.*

Mycobank no.: MB 501299

Basidiomata resupinata, effuso-reflexa vel discoidea; hymenophora laevia; systema hypharum monomiticum; dendrohyphidia raro absentia; basidia saepe e probasidiis oriuntur. Cystidia presentia vel absentia. Sporae hyalinae, tenuitunicatae, albae vel aggregatae roseae.

Typus: *Corticium* Pers. 1794.

Basidiomycetes with effused or discoid (*Cyrtidia*) basidiomata, a smooth hymenophore, and a monomitic hyphal system with clamped, rarely simple-septate, hyphae. Dendrohyphidia common. Species with or without cystidia. A probasidial resting stage is present in many species. Spores smooth, in masses white to pink. Saprotrophic, parasitic, or lichenicolous.

Equivalent to *Vuilleminiales* Boidin et al. 1998 and the corticioid clade (Binder et al. 2005; Larsson et al. 2004). Boidin et al. (1998) explicitly included *Corticium* in their new order, as a member of the family *Vuilleminiaceae* Maire 1902. Jülich (1981) also placed *Corticium* in *Vuilleminiaceae* but referred them to *Aleurodiscales* Jülich 1981. *Corticium* is the type of *Corticaceae* Herter 1910, a family name conserved against *Vuilleminiaceae*. The introduction of *Corticiales* as a new name for this order is, therefore, the preferred option.

Exemplar genera: *Corticium* Pers. 1794, *Vuilleminia* Maire 1902, *Punctularia* Pat. 1895.

Order: **Gloeophyllales** Thorn, *ord. nov.*

Mycobank no.: MB 501300

Basidiomata annua vel perennia, resupinata, effuso-reflexa, dimidiata vel pileata; hymenophora laevia, merulioidea, odontioidea vel poroidea. Systema hypharum monomiticum, dimiticum vel trimiticum. Hyphae generativae fibulatae vel efibulatae. Leptocystidia ex trama in hymenium projecta, hyalina vel brunnea, tenuitunicata vel crassitunicata. Basidiosporae laeves, hyalinae, tenuitunicatae, ellipsoideae vel cylindricae vel allantoideae, inamyloideae. Lignum decompositum brunneum vel album.

Typus: *Gloeophyllum* P. Karst. 1882.

Fruiting bodies perennial or annual and long-lived, with hymenium maturing and thickening over time. Stature resupinate, effused-reflexed or dimidiate, with smooth, wrinkled,

dentate, lamellate or regularly poroid hymenophore, or pileate-stipitate with lamellae. (Aborted, coralloid or flabelliform fruiting bodies may be formed under conditions of darkness or high carbon dioxide concentration). Leptocystidia or hyphoid hairs originating in the context and extending into or protruding from the hymenial layer (or lamellar margin in *Neolentinus*) are common; these often with thick brown walls and brownish incrustation. Context brown (but pallid in *Neolentinus*) and generally darkening in potassium hydroxide (the brownish incrustation in *Boreostereum* turning green in potassium hydroxide). Monomitic (if so, with sclerified generative hyphae), dimitic, or trimitic; generative hyphae with or without clamp connections. Basidiospores hyaline, ellipsoid to cylindrical or suballantoid, with thin, smooth walls, and neither amyloid, dextrinoid nor cyanophilous. Where this is known, basidiospores are binucleate and sexuality is heterothallic and bipolar (but tetrapolar in *V. berkeleyi*).

Causing brown rots (*Gloeophyllum*, *Neolentinus*, *Veluticeps*) or stringy white rot (*Boreostereum*, *Donkioporia*) of wood of gymnosperms, monocots and dicots. Occurrence on 'wood in service' (e.g. railway ties, paving blocks, wooden chests) seems to be common (in *Donkioporia*, *Gloeophyllum*, *Heliocybe* and *Neolentinus*); often on charred wood (*Boreostereum* and *Veluticeps*).

Equivalent to *Gloeophyllum* clade (Binder et al. 2005).

Exemplar genera: *Gloeophyllum* P. Karst. 1882, *Neolentinus* Redhead & Ginns 1985, *Veluticeps* (Cooke) Pat. 1894.

Order: **Hymenochaetales** Oberw., in Frey et al. (eds), *Beitr. Biol. niederen Pflanz.*: 89 (1977).

Equivalent to the hymenochaetoid clade (Hibbett & Thorn 2001; Larsson et al. 2007).

Exemplar genera: *Hymenochaete* Lév. 1846, *Phellinus* Quél. 1886, *Trichaptum* Murrill 1904.

Order: **Polyporales** Gäum., *Vergl. Morph. Pilze*: 503 (1926).

Equivalent to polyporoid clade (Hibbett & Thorn 2001).

Exemplar genera: *Polyporus* Fr. 1815, *Fomitopsis* P. Karst. 1881, *Phanerochaete* P. Karst. 1889.

Order: **Russulales** Kreisel ex P. M. Kirk, P. F. Cannon & J. C. David, in Kirk et al., *Ainsworth & Bisby's Dict. Fungi* (9th edn): xi (2001).

Equivalent to the russuloid clade (Hibbett & Thorn 2001; Larsson & Larsson 2003; Miller et al. 2007).

Exemplar genera: *Russula* Pers. 1796, *Aleurodiscus* Rabenh. ex J. Schröt. 1888, *Bondarzewia* Singer 1940, *Hericium* Pers. 1794, *Peniophora* Cooke 1879, *Stereum* Pers. 1794.

Order: **Sebacinales** M. Weiß, Selosse, Rexer, A. Urb. & Oberw., *Mycol. Res.* 108: 1007 (2004b).

Exemplar genera: *Sebacina* Tul. 1871, *Tremellodendron* G. F. Atk. 1902, *Piriformospora* Sav. Verma, Aj. Varma, Rexer, G. Kost & P. Franken 1998.

Order: **Thelephorales** Corner ex Oberw., *Sydowia* 78: 361 (1976).

Equivalent to the thelephoroid clade (Hibbett & Thorn 2001).

Exemplar genera: *Thelephora* Ehrh. ex Willd. 1787, *Bankera* Coker & Beers ex Pouzar 1955, *Polyozellus* Murrill 1910.

Order: **Trechisporales** K. H. Larss., ord. nov.

MycoBank no.: MB 501301

Basidiomata resupinata, stipitata vel clavarioidea. Hymenophora laevia, grandinioidea, hydnoidea vel poroidea. Systema hypharum monomiticum vel dimiticum. Hyphae fibulatae, septa hypharum interdum inflata (ampullata). Cystidia praesentia vel absentia. Basidia 4-6 sterigmata formantia. Sporae laeves vel ornatae. Species lignicolae vel terricolae.

Typus: *Trechispora* P. Karst. 1890.

Basidiomycetes with effused, stipitate or clavarioid basidiomata. Hymenophore smooth, grandinoid, hydroid or poroid. Hyphal system monomitic, hyphae clamped, subcircular hyphae with or without ampullate septa. Cystidia present in some species, mostly lacking. Basidia with four to six sterigmata. Spores smooth or ornamented. On wood or soil.

Equivalent to *Hydnodontales* Jülich 1981 and trechisporoid clade (Binder et al. 2005; Larsson et al. 2004). *Hydnodon* Banker 1913 was recently placed in synonymy under *Trechispora* (Ryvarden 2002) and this synonymy is supported by molecular data (K.H. Larsson, unpubl.). The introduction of a new name for the group, a name that connects to the clade name already established and that is based on the most species-rich genus is, therefore, justified.

Exemplar genera: *Trechispora* P. Karst. 1890, *Sistotremastrum* J. Erikss. 1958, *Porpomyces* Jülich 1982.

Basidiomycota incertae sedis (not placed in any subphylum):

Class: **Wallemiomycetes** Zalar, de Hoog & Schroers, *Antonie van Leeuwenhoek* 87: 322 (2005).

Analyses of *rpb1*, *rpb2*, *tef1*, and *nuc-lsu*, *nuc-ssu*, and 5.8S ribosomal RNA genes suggest that the *Wallemiomycetes* is the sister group of the rest of the *Basidiomycota* (possibly along with *Entorrhizomycetes*, see below), but subsets of this dataset produce alternative placements (Matheny et al. 2007b; Zalar et al. 2005).

Order: **Wallemiales** Zalar, de Hoog & Schroers, *Antonie van Leeuwenhoek* 87: 322 (2005).

Exemplar genus: *Wallemia* Johan-Olsen 1887.

Class: **Entorrhizomycetes** Begerow, Stoll & R. Bauer, *Mycologia* 98: 908 (2007) ['2006'].

Equivalent to *Entorrhizomycetidae* R. Bauer & Oberw. (Bauer et al. 1997). So far, only ribosomal RNA genes have been sequenced in *Entorrhizomycetes*. Analyses with broad sampling across all groups of *Basidiomycota* and including *Ascomycota* and *Glomeromycota* as outgroups suggest that *Entorrhizomycetes* is not nested within any subphylum, and may be the sister group of the rest of the *Basidiomycota* (Matheny et al. 2007a; also see Begerow et al. 1997).

Order: **Entorrhizales** R. Bauer & Oberw., in Bauer et al., *Can. J. Bot.* 75: 1311 (1997).

Exemplar genus: *Entorrhiza* C. A. Weber 1884.

Acknowledgments

Preparation of this classification was facilitated by the Assembling the Fungal Tree of Life (AFTOL) project and the Deep Hypha Research Coordination Network, which were supported by the US National Science Foundation grants DEB-0228657, DEB-0228668, DEB-0228671, DEB-0228725 and NSF-0090301.

Note added in proof

After this article went to press, the authors became aware of the following publication, which includes alternative citations for many of the names included here: Doweld A, 2001. *Prosyllabus tracheophytorum: Tentamen systematis plantarum vascularium (Tracheophyta)*. Geos, Moscow.

REFERENCES*

- Adl SM, Simpson AGB, Farmer MA, Andersen RA, Anderson OR, Barta JR, Bowser SS, Brugerolle G, Fensome RA, Fredericq S, James TY, Karpov S, Kugrens P, Krug J, Lane CE, Lewis LA, Lodge J, Lynn DH, Mann DG, McCourt RM, Mendoza L, Moestrup Ø, Mozley-Standridge SE, Nerad TA, Shearer CA, Smirnov AV, Spiegel FW, Taylor MFJR, 2005. The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. *Journal of Eukaryotic Microbiology* 52: 399–451.
- Aime MC, Matheny PB, Henk DA, Frieders EM, Nilsson RH, Piepenbring M, McLaughlin DJ, Szabo LJ, Begerow D, Sampaio JP, Bauer R, Weiß M, Oberwinkler F, Hibbett DS, 2007 ['2006']. An overview of the higher-level classification of *Pucciniomycotina* based on combined analyses of nuclear large and small subunit rDNA sequences. *Mycologia* 98: 896–905.
- Alexopoulos CJ, Mims CW, Blackwell M, 1996. *Introductory Mycology*, fourth ed. John Wiley, New York.
- von Arx JA, 1967. *Pilzkunde*. J. Cramer, Lehre.
- Balbani G, 1882. Sur les microsporides ou psorospermies des Articulés. *Comptes rendus de l'Académie des Sciences, Paris* 95: 1168–1171.
- Baldauf SL, Roger AJ, Wenk-Siefert I, Doolittle WF, 2000. A kingdom-level phylogeny of eukaryotes based on combined protein data. *Science* 290: 972–977.
- Barr DJS, 1980. An outline for the reclassification of the *Chytriales*, and for a new order, the *Spizellomycetales*. *Canadian Journal of Botany* 58: 2380–2394.
- Barr ME, 1983. The ascomycete connection. *Mycologia* 75: 1–13.
- Barr ME, 1987a. New taxa and combinations in the *Loculoascomycetes*. *Mycotaxon* 29: 501–505.
- Barr ME, 1987b. *Prodromus to Class Loculoascomycetes*. University of Massachusetts, Amherst.
- Barron GL, 1975. Nematophagus fungi: *Helicocephalum*. *Transactions of the British Mycological Society* 65: 309–310.
- Bauer R, Begerow D, Oberwinkler F, Piepenbring M, Berbee ML, 2001. *Ustilaginomycetes*. In: McLaughlin DJ, McLaughlin EJ, Lemke P (eds), *The Mycota*. Vol. VIII: Part B. *Systematics and Evolution*. Springer-Verlag, Berlin, pp. 57–84.
- Bauer R, Begerow D, Sampaio JP, Weiß M, Oberwinkler F, 2006. The simple-septate basidiomycetes: a synopsis. *Mycological Progress* 5: 41–66.
- Bauer R, Oberwinkler F, Vánky K, 1997. Ultrastructural markers and systematics in smut fungi and allied taxa. *Canadian Journal of Botany* 75: 1273–1314.
- Begerow D, Bauer R, Boekhout T, 2000. Phylogenetic placements of ustilaginomycetous anamorphs as deduced from nuclear LSU rDNA sequences. *Mycological Research* 104: 53–60.
- Begerow D, Bauer R, Oberwinkler F, 1997. Phylogenetic studies on nuclear large subunit ribosomal DNA sequences of smut fungi and related taxa. *Canadian Journal of Botany* 75: 2045–2056.
- Begerow D, Stoll M, Bauer R, 2007 ['2006']. A phylogenetic hypothesis of *Ustilaginomycotina* based on multiple gene analyses and morphological data. *Mycologia* 98: 906–916.
- Benjamin RK, 1979. *Zygomycetes and their spores*. In: Kendrick B (ed), *The Whole Fungus: the Sexual–Asexual Synthesis*. National Museum of Natural Sciences, Ottawa, pp. 573–616.
- Benny GL, Kimbrough JW, 1980. A synopsis of the orders and families of *Plectomycetes* with keys to genera. *Mycotaxon* 12: 1–91.
- Benny GL, White MM, 2001. The classification and phylogeny of *Trichomycetes* and *Zygomycetes*. In: Misra JK, Horn BW (eds), *Trichomycetes and Other Fungal Groups*. Science Publishers, Enfield, NH, pp. 39–53.
- Berres ME, Szabo LJ, McLaughlin DJ, 1995. Phylogenetic relationships in auriculariaceous basidiomycetes based on 25S ribosomal DNA sequences. *Mycologia* 87: 821–840.
- Binder M, Hibbett DS, Larsson K-H, Larsson E, Langer E, Langer G, 2005. The phylogenetic distribution of resupinate forms across the major clades of mushroom-forming fungi (*Homobasidiomycetes*). *Systematics and Biodiversity* 3: 1–45.
- Binder M, Hibbett DS, 2007 ['2006']. Molecular systematics and biological diversification of *Boletales*. *Mycologia* 98: 971–983.
- Blackwell M, Hibbett DS, Taylor JW, Spatafora JW, 2007 ['2006']. Research coordination networks: a phylogeny for kingdom *Fungi*. *Mycologia* 98: 829–837.
- Boidin J, Mugnier J, Canales R, 1998. Taxonomie moléculaire des *Aphyllphorales*. *Mycotaxon* 66: 445–492.
- Bold HC, 1957. *The Morphology of Plants*. Harper Row, New York.
- Bullerwell CE, Forget L, Lang BF, 2003. Evolution of monoblepharidalean fungi based on complete mitochondrial genome sequences. *Nucleic Acids Research* 31: 1614–1623.
- Campbell J, Anderson JL, Shearer CA, 2003. Systematics of *Halosarphaea* based on morphological and molecular data. *Mycologia* 95: 530–552.
- Campbell J, Volkmann-Kohlmeyer B, Gräfenhan T, Spatafora JW, Kohlmeyer J, 2005. A re-evaluation of *Lulworthiales*: relationships based on 18S and 28S rDNA. *Mycological Research* 109: 556–568.
- Castlebury LA, Rossman AY, Jaklitsch WJ, Vasilyeva LN, 2002. A preliminary overview of the *Diaportheales* based on large subunit nuclear ribosomal DNA sequences. *Mycologia* 94: 1017–1031.
- Castlebury LA, Rossman AY, Sung G-H, Hyten AS, Spatafora JW, 2004. Multigene phylogeny reveals new lineage for *Stachybotrys chartarum*, the indoor air fungus. *Mycological Research* 108: 864–872.
- Cavalier-Smith T, 1981. Eukaryote kingdoms: seven or nine? *BioSystems* 14: 461–481.
- Cavalier-Smith T, 1998. A revised six-kingdom system of Life. *Biological Reviews* 73: 203–266.
- Cejp K, 1957. *Houby; celostátní vysokoškolská učebnice*. Vol. 1. Czechoslovakian Academy of Sciences Press, Prague.
- Clements FE, Shear CL, 1931. *The Genera of Fungi*, second ed. H.W. Wilson, New York.
- Cohn F, 1879. Ueber ein Thallophytensystem. *Jahresbericht der Schlesischen Gesellschaft für vaterländische Cultur, Breslau* 57: 279.
- Cracraft J, Donoghue MJ (eds), 2004. *Assembling the Tree of Life*. Oxford University Press, Oxford.
- Currah RS, 1985. Taxonomy of the *Onygenales*: *Arthrodermataceae*, *Gymnoaceae*, *Myxotrichaceae* and *Onygenaceae*. *Mycotaxon* 24: 1–216.

* This list of references, in addition to including papers cited in the text, also contains the full bibliographic details of some papers otherwise cited only as places of publications of names where those works may otherwise be difficult to locate.

- David JC, 2002. A preliminary catalogue of the names of fungi above the rank of order. *Constancea* **83**: 1–30.
- Del Prado R, Schmitt I, Kautz S, Palice Z, Luecking R, Lumbsch HT, 2006. Molecular data place *Trypetheliaceae* in *Dothideomycetes*. *Mycological Research* **110**: 511–520.
- Donk MA, 1964. A conspectus of the families of *Aphyllophorales*. *Persoonia* **3**: 199–324.
- Ekman S, Tønnsberg T, 2002. Most species of *Lepraria* and *Lepruloma* form a monophyletic group closely related to *Stereocaulon*. *Mycological Research* **106**: 1262–1276.
- Engler A, 1898. *Syllabus der Pflanzenfamilien: Eine Übersicht über das gesammte Pflanzensystem*, second ed. Gebrüder Borntraeger Verlag, Berlin, 46.
- Eriksson O, 1982. Outline of the ascomycetes. *Mycotaxon* **15**: 203–248.
- Eriksson OE, 1986. *Lahmia* Körber (= *Parkerella* A. Funk) a misinterpreted genus with isolated position. *Mycotaxon* **27**: 347–360.
- Eriksson OE, 1992. *Huangshania verrucosa* gen. et sp. nov. (Triblidaceae, Triblidiales ordo nov.), a discomycete on *Pinus* from China. *Systema Ascomycetum* **11**: 1–10.
- Eriksson OE, 1994. *Pneumocystis carinii*, a parasite in lungs of mammals, referred to a new family and order (*Pneumocystidaceae*, *Pneumocystidales*, *Ascomycota*). *Systema Ascomycetum* **13**: 165–180.
- Eriksson OE, Baral H-O, Currah RS, Hansen K, Kurtzman CP, Rambold G, Læssøe T, 2003. Notes on ascomycete systematics. Nos 3580–3623. *Myconet* **9**: 91–103.
- Eriksson OE, Svedskog A, Landvik S, 1993. Molecular evidence for the evolutionary hiatus between *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*. *Systema Ascomycetum* **11**: 119–162.
- Eriksson OE, Winka K, 1997. Supraordinal taxa of *Ascomycota*. *Myconet* **1**: 1–16.
- Fell JW, Boekhout T, Fonseca A, Sampaio JP, 2001. Basidiomycetous yeasts. In: McLaughlin DJ, McLaughlin EJ, Lemke P (eds), *The Mycota*. Vol. VII. Part B., *Systematics and Evolution*. Springer-Verlag, Berlin, pp. 3–35.
- Fell JW, Roeijmans H, Boekhout T, 1999. A new order of basidiomycetous yeasts. *International Journal of Systematic Bacteriology*. *Cystofilobasidiales* **49**: 907–913.
- Fell JW, Scorzetti G, 2004. Reassignment of the basidiomycetous yeasts *Trichosporon pullulans* to *Guehomyces pullulans* gen. nov., comb. nov. and *Hyalodendron lignicola* to *Trichosporon lignicola* comb. nov. *International Journal of Systematic and Evolutionary Microbiology* **54**: 995–998.
- Fischer A, 1892. *Phycomycetes. Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz* **1**: 1–128.
- Fischer E, 1898. Phallineae. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien nebst ihren Gattungen und wichtigeren Arten insbesondere der Nutzpflanzen*, Vol. 1. Wilhelm Engelmann, Leipzig, pp. 276–295.
- Fries EM, 1821. *Systema Mycologicum*, Vol. 1. Berling, Lund.
- Fries EM, 1832. *Systema Mycologicum*, Vol. 3. Ernest Mauriti, Greifswald.
- Galagan JE, Henn MR, Ma L-J, Cuomo CA, Birren B, 2005. Genomics of the fungal kingdom: insights into eukaryotic biology. *Genome Research* **15**: 1620–1631.
- Gamundí IJ, 1971. Las *Cyrtariales* sudamericanas. *Darwiniana* **16**: 461–510.
- Gäumann E, 1926. *Vergleichende Morphologie der Pilze*. Gustav Fischer, Jena.
- Gäumann E, Dodge CW, 1928. *Comparative Morphology of the Fungi*. McGraw-Hill, London.
- Geiser DM, Gueidan C, Miądlikowska J, Lutzoni F, Kauff F, Hofstetter V, Fraker E, Schoch CL, Tibell L, Untereiner WA, Aptroot A, 2007 [‘2006’]. *Eurotiomycetes: Eurotiomycetidae* and *Chaetothyriomycetidae*. *Mycologia* **98**: 1054–1065.
- Germot A, Philipe H, Le Guyader H, 1997. Evidence for loss of mitochondria in microsporidia from a mitochondrial HSP70 in *Nosema locustae*. *Molecular and Biochemical Parasitology* **87**: 159–168.
- Gibson JL, Kimbrough JW, Benny GL, 1986. Ultrastructural observations on *Endogonaceae* (*Zygomycetes*). II. *Glaziellales* ord. nov. and *Glaziellaceae* fam. nov.: new taxa based upon light and electron microscopic observations of *Glaziella aurantiaca*. *Mycologia* **78**: 941–954.
- Gilbert EJ, 1931. *Les Bolets*. [Les Livres du Mycologue, Vol. 3]. E. Le François, Paris.
- Gill EE, Fast NM, 2006. Assessing the microsporidia–fungi relationship: combined phylogenetic analysis of eight genes. *Gene* **375**: 103–109.
- Grube M, Baloch E, Lumbsch HT, 2004. The phylogeny of *Porinaceae* (*Ostropomycetidae*) suggests a neotenic origin of perithecia in *Lecanoromycetes*. *Mycological Research* **108**: 1111–1118.
- Gwynne-Vaughan HCI, 1922. *Fungi: Ascomycetes, Ustilaginales, Uredinales*. Cambridge University Press, Cambridge.
- Hausner G, Reid J, 2004. The nuclear small subunit ribosomal genes of *Sphaeronaemella helvellae*, *Sphaeronaemella fimicola*, *Gabarnaudia betae*, and *Cornuvesica falcata*: phylogenetic implications. *Canadian Journal of Botany* **82**: 752–762.
- Hawksworth DL, Kirk PM, Sutton BC, Pegler DN, 1995. *Ainsworth & Bisby's Dictionary of the Fungi*, eighth ed. CAB International, Wallingford.
- Henk DA, Weir A, Blackwell M, 2003. *Laboulbeniopsis*, an ectoparasite of termites newly recognized as a member of the *Laboulbeniomyces*. *Mycologia* **95**: 561–564.
- Hennings P, 1897. *Exobasidiineae*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 103–105.
- Hennings P, 1898. *Dacryomycetinae*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 96–102.
- Henssen A, Jahms HM, 1973 [1974]. *Lichenes: Eine Einführung in die Flechtenkunde*. George Thieme, Stuttgart.
- Hibbett DS, 2007 [‘2006’]. A phylogenetic overview of the *Agaricomycotina*. *Mycologia* **98**: 917–925.
- Hibbett DS, Donoghue MJ, 1998. Integrating phylogenetic analysis and classification in fungi. *Mycologia* **90**: 347–356.
- Hibbett DS, Thorn RG, 2001. *Homobasidiomycetes*. In: McLaughlin DJ, McLaughlin EJ, Lemke P (eds), *The Mycota*. Vol. VII. Part B., *Systematics and Evolution*. Springer-Verlag, Berlin, pp. 121–168.
- Hirt RP, Healy B, Vossbrinck CR, Canning EU, Embley TM, 1997. A mitochondrial Hsp70 orthologue in *Vairimorpha necatrix*: molecular evidence that Microsporidia once contained mitochondria. *Current Biology* **7**: 995–998.
- Hofstetter V, Miądlikowska J, Kauff F, Lutzoni F, 2007. Phylogenetic comparison of protein-coding versus ribosomal RNA-coding sequence data: a case study of the *Lecanoromycetes* (*Ascomycota*). *Molecular Phylogenetics and Evolution*, in press.
- de Hoog GS, Göttlich E, Platas G, Genilloud O, Leotta G, Brummelen J, 2005. Evolution, taxonomy and ecology of the genus *Thelebolus* in Antarctica. *Studies in Mycology* **51**: 33–76.
- Hosaka K, Bates ST, Beever RT, Castellano MA, Colgan III W, Domínguez LS, Nouhra ER, Geml J, Giachini AJ, Kenney SR, Simpson NB, Spatafora JW, Trappe JM, 2007 [‘2006’]. Molecular phylogenetics of the gomphoid-phalloid fungi with an establishment of the new subclass *Phallomycetidae* and two new orders. *Mycologia* **98**: 949–959.
- Huhndorf SM, Miller AN, Fernández FA, 2004a. Molecular systematics of the *Coronophorales* and new species of

- Bertia*, *Lasiobertia* and *Nitschkia*. *Mycological Research* **108**: 1384–1398.
- Huhndorf SM, Miller AN, Fernández FA, 2004b. Molecular systematics of the *Sordariales*: the order and the family *Lasiosphaeriaceae* redefined. *Mycologia* **96**: 368–387.
- Inderbitzin P, Landvik S, Abdel-Wahab MA, Berbee ML, 2001. *Aliquandostipitaceae*, a new family for two new tropical ascomycetes with unusually wide hyphae and dimorphic ascospores. *American Journal of Botany* **88**: 52–61.
- Inderbitzin P, Lim SR, Volkmann-Kohlmeyer B, Kohlmeyer J, Berbee ML, 2004. The phylogenetic position of *Spathulospora* based on DNA sequences from dried herbarium material. *Mycological Research* **108**: 737–748.
- International Commission on Zoological Nomenclature, 1999. *International Code of Zoological Nomenclature*. International Trust for Zoological Nomenclature, London.
- Jahn TL, Jahn FF, 1949. *How to Know the Protozoa*. Wm C. Brown, Dubuque.
- James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, Cox C, Celio G, Gueidan C, Fraker E, Miądlikowska J, Lumbsch HT, Rauhut A, Reeb V, Arnold EA, Amtoft A, Stajich JE, Hosaka K, Sung G-H, Johnson D, O'Rourke B, Crockett M, Binder M, Curtis JM, Slot JC, Wang Z, Wilson AW, Schüßler A, Longcore JE, O'Donnell K, Mozley-Standridge S, Porter D, Letcher PM, Powell MJ, Taylor JW, White MM, Griffith GW, Davies DR, Humber RA, Morton J, Sugiyama J, Rossman AY, Rogers JD, Pfister DH, Hewitt D, Hansen K, Hambleton S, Shoemaker RA, Kohlmeyer J, Volkmann-Kohlmeyer B, Spotts RA, Serdani M, Crous PW, Hughes KW, Matsuura K, Langer E, Langer G, Untereiner WA, Lücking R, Büdel B, Geiser DM, Aptroot A, Diederich P, Schmitt I, Schultz M, Yahr R, Hibbett DS, Lutzoni F, McLaughlin D, Spatafora J, Vilgalys R, 2006. Reconstructing the early evolution of the fungi using a six gene phylogeny. *Nature* **443**: 818–822.
- James TY, Letcher PM, Longcore JE, Mozley-Standridge SE, Porter D, Powell MJ, Griffith GW, Vilgalys R, 2007 [‘2006’]. A molecular phylogeny of the flagellated fungi (*Chytridiomycota*) and a proposal for a new phylum (*Blastocladiomycota*). *Mycologia* **98**: 860–871.
- James TY, Porter D, Leander CA, Vilgalys R, Longcore JE, 2000. Molecular phylogenetics of the *Chytridiomycota* supports the utility of ultrastructural data in chytrid systematics. *Canadian Journal of Botany* **78**: 336–350.
- Jülich W, 1981. Higher taxa of basidiomycetes. *Bibliotheca Mycologica* **85**: 1–485.
- Kauff F, Lutzoni F, 2002. Phylogeny of the *Gyalectales* and *Ostropales* (*Ascomycota*, *Fungi*): among and within order relationships based on nuclear ribosomal RNA small and large subunits. *Molecular Phylogenetics and Evolution* **25**: 138–156.
- Keeling PJ, 2003. Congruent evidence for alpha-tubulin and beta-tubulin gene phylogenies for a zygomycete origin of *Microsporidia*. *Fungal Genetics and Biology* **38**: 298–309.
- Keeling PJ, Luker MA, Palmer JD, 2000. Evidence from beta-tubulin phylogeny that *Microsporidia* evolved from within the fungi. *Molecular Biology and Evolution* **17**: 23–31.
- Kendrick B (ed), 1979. *The Whole Fungus: the Sexual-asexual Synthesis*, Vol. 2. National Museum of Natural Sciences, Ottawa.
- Kendrick B, 1985. *The Fifth Kingdom*. Mycologue Publications, Waterloo.
- Kirk PM, Cannon PF, David JC, Stalpers JA (eds), 2001. *Ainsworth & Bisby's Dictionary of the Fungi*, ninth ed. CABI Publishing, Wallingford.
- Kohlmeyer J, Spatafora JW, Volkmann-Kohlmeyer B, 2000. *Lulworthiales*, a new order of marine *Ascomycota*. *Mycologia* **92**: 453–458.
- Kruys A, Eriksson OE, Wedin M, 2006. Phylogenetic relationships of coprophilous *Pleosporales* (*Dothideomycetes*, *Ascomycota*), and the classification of some bitunicate taxa of unknown position. *Mycological Research* **110**: 527–536.
- Kuramae EE, Robert V, Snel B, Weiß M, Boekhout T, 2006. Phylogenomics reveal a robust fungal tree of life. *FEMS Yeast Research* **6**: 1213–1220.
- Kudrjavec W, 1960. *Die Systematik der Hefen*. Akademie Verlag, Berlin.
- Kurtzman CP, Sugiyama J, 2001. Ascomycetous yeasts and yeastlike taxa. In: McLaughlin DJ, McLaughlin EJ, Lemke P (eds), *The Mycota*. Vol. VII. Part A. *Systematics and Evolution*. Springer-Verlag, Berlin, pp. 179–200.
- Landvik S, Eriksson OE, 1994. Relationship of the genus *Glaziella* (*Ascomycota*) inferred from 18S rDNA sequences. *Systema Ascomycetum* **13**: 13–23.
- Landvik S, Eriksson OE, Berbee ML, 2001. *Neolecta* — a fungal dinosaur? Evidence from β -tubulin amino acid sequences. *Mycologia* **93**: 1151–1163.
- Landvik S, Eriksson OE, Gargas A, Gustafsson P, 1993. Relationships of the genus *Neolecta* (*Neolectales* ordo nov., *Ascomycotina*) inferred from 18S rDNA sequences. *Systema Ascomycetum* **11**: 107–118.
- Larsson E, Larsson K-H, 2003. Phylogenetic relationships of rust-like basidiomycetes with emphasis on aphyllorphorean taxa. *Mycologia* **95**: 1037–1065.
- Larsson JIR, 2000. The hyperparasitic microsporidium *Amphiacantha longa* Caullery et Mesnil, 1914 (*Microspora: Metchnikovellidae*) — description of the cytology, redescription of the species, emended diagnosis of the genus *Amphiacantha* and establishment of the new family *Amphiacanthidae*. *Folia Parasitologica* **47**: 241–256.
- Larsson K-H, Larsson E, Kõljalg U, 2004. High phylogenetic diversity among corticioid homobasidiomycetes. *Mycological Research* **108**: 983–1002.
- Larsson K-H, Parmasto E, Fischer M, Langer E, Nakasone KK, Redhead SA, 2007 [‘2006’]. *Hymenochaetales*: a molecular phylogeny for the hymenochaetoid clade. *Mycologia* **98**: 926–936.
- Letcher PM, Powell MJ, Churchill PF, Chambers JG, 2006. Ultrastructural and molecular phylogenetic delineation of a new order, the *Rhizophydiales* (*Chytridiomycota*). *Mycological Research* **110**: 898–915.
- Li JL, Heath IB, Packer L, 1993. The phylogenetic relationships of the anaerobic chytridiomycetous gut fungi (*Neocallimastaceae*) and the *Chytridiomycota*. II. Cladistic analysis of structural data and description of *Neocallimasticales* ord. nov. *Canadian Journal of Botany* **71**: 393–407.
- Lichtwardt RW, Manier JF, 1978. Validation of the *Harpellales* and *Asellariales*. *Mycotaxon* **7**: 441–442.
- Lindau G, 1896. *Hysteriineae*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 265–278.
- Lindau G, 1897a. *Dothideales*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 373–383.
- Lindau G, 1897b. *Hypocreales*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1(1). Leipzig. Wilhelm Engelmann Verlag, pp. 343–372.
- Lindau G, 1897c. *Laboulbeniineae*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1(1). Leipzig. Wilhelm Engelmann Verlag, pp. 491–505.
- Lindemuth R, Wirtz N, Lumbsch HT, 2001. Phylogenetic analysis of nuclear and mitochondrial rDNA sequences supports the view that loculoascomycetes (*Ascomycota*) are not monophyletic. *Mycological Research* **105**: 1176–1181.
- Liu YJ, Hall BD, 2004. Body plan evolution of ascomycetes, as inferred from an RNA polymerase II phylogeny. *Proceedings of the National Academy of Sciences, USA* **101**: 4507–4512.

- Liu YJ, Hodson MC, Hall BD, 2006. Loss of the flagellum happened only once in the fungal lineage: phylogenetic structure of kingdom Fungi inferred from RNA polymerase II subunit genes. *BMC Evolutionary Biology* 6: 74. www.biomedcentral.com/1471-2148/6/74.
- Locquin M, 1984. *Mycologie Générale et Structurale*. Masson, Paris.
- Lücking R, Stuart B, Lumbsch HT, 2004. Phylogenetic relationships of Gomphillaceae and Asterothyriaceae: evidence from a combined Bayesian analysis of nuclear and mitochondrial sequences. *Mycologia* 96: 283–294.
- Lumbsch T, Palice Z, Wiklund E, Ekman S, Wedin M, 2004. Supraordinal phylogenetic relationships of Lecanoromycetes based on Bayesian analysis of combined nuclear and mitochondrial sequences. *Molecular Phylogenetics and Evolution* 31: 822–832.
- Lumbsch HT, Schmitt I, Lindemuth R, Miller A, Mangold A, Fernandez F, Huhndorf S, 2005. Performance of four ribosomal DNA regions to infer higher-level phylogenetic relationships of inoperculate euascomycetes (Leotiomyceta). *Molecular Phylogenetics and Evolution* 34: 512–524.
- Lumbsch HT, Schmitt I, Lücking R, Wiklund RE, Wedin M, 2007. The phylogenetic placement of Ostropales within Lecanoromycetes (Ascomycota) revisited. *Mycological Research* 111.
- Lumbsch HT, Wirtz N, Lindemuth R, Schmitt I, 2002. Higher level phylogenetic relationships of euascomycetes (Pezizomycotina) inferred from a combined analysis of nuclear and mitochondrial sequence data. *Mycological Progress* 1: 57–70.
- Lutzoni F, Kauff F, Cox CJ, McLaughlin D, Celio G, Dentinger B, Padamsee M, Hibbett DS, James TY, Baloch E, Grube M, Reeb V, Hofstetter V, Schoch C, Arnold AE, Miądlikowska J, Spatafora J, Johnson D, Hambleton S, Crockett M, Shoemaker R, Sung G-H, Lücking R, Lumbsch T, O'Donnell K, Binder M, Diederich P, Ertz D, Gueidan C, Hansen K, Harris RC, Hosaka K, Lim Y-W, Matheny B, Nishida H, Pfister D, Rogers J, Rossmann A, Schmitt I, Sipman H, Stone J, Sugiyama J, Yahr R, Vilgalys R, 2004. Assembling the fungal tree of life: progress, classification, and evolution of subcellular traits. *American Journal of Botany* 91: 1446–1480.
- Manier J-F, 1973. Quelques aspects ultrastructuraux du trichomycète asellariale, *Asellaria ligae* Tuzet et Manier, 1950 ex Manier, 1958. *Comptes Rendu hebdomadaires des séances de l'Académie des Sciences de Paris, séries D* 276: 3429–3431.
- Margulis L, Corliss JO, Melkonian M, Chapman DJ, 1990. *Handbook of Protoctista: the structure, cultivation, habitats and life histories of the eukaryotic micro-organisms and their descendants*. Jones & Bartlett, Boston.
- Matheny PB, Curtis JM, Hofstetter V, Aime MC, Moncalvo J-M, Ge Z-W, Yang Z-L, Slot JC, Ammirati JF, Baroni TJ, Bougher NL, Hughes KW, Lodge DJ, Kerrigan RW, Seidl MT, Aanen DK, DeNitis M, Daniele GM, Desjardin DE, Kropp BR, Norvell LL, Parker A, Vellinga EC, Vilgalys R, Hibbett DS, 2007 ["2006"]a. Major clades of Agaricales: a multi-locus phylogenetic overview. *Mycologia* 98: 984–997.
- Matheny PB, Gossman JA, Zalar P, Arun Kumar TK, Hibbett DS, 2007a. Resolving the phylogenetic position of the Wallemiomycetes: an enigmatic major lineage of Basidiomycota. *Canadian Journal of Botany* 84: 1794–1805.
- Matheny PB, Wang Z, Binder M, Curtis JM, Lim YW, Nilsson RH, Hughes KW, Hofstetter V, Ammirati JF, Schoch CL, Langer GE, McLaughlin DJ, Wilson AW, Frøslev T, Ge ZW, Kerrigan RW, Slot JC, Vellinga EC, Liang ZL, Baroni TJ, Fischer M, Hosaka K, Matsuura K, Seidl MT, Vauria J, Hibbett DS, 2006. Contributions of *rpb2* and *tef1* to the phylogeny of mushrooms and allies (Basidiomycota, Fungi). *Molecular Phylogenetics and Evolution* in press.
- McLaughlin DJ, McLaughlin EG, Lemke PA (eds), 2001a. *The Mycota*. Vol. VII. Part A. Systematics and Evolution. Springer-Verlag, Berlin.
- McLaughlin DJ, McLaughlin EG, Lemke PA (eds), 2001b. *The Mycota*. Vol. VII. Part B. Systematics and Evolution. Springer-Verlag, Berlin.
- McNeill JF, Barrie F, Burdet HM, Demoulin V, Hawksworth DL, Marhold K, Nicolson DH, Prado J, Silva PC, Skog JE, Wiersema J, Turland NJ (eds), 2006. *International Code of Botanical Nomenclature (Vienna Code) [Regnum Vegetabile Vol. 146]*. A.R.G. Ganter Verlag, Ruggell.
- Miądlikowska J, Kauff F, Hofstetter V, Fraker E, Reeb V, Grube M, Hafellner J, Kukwa M, Lücking R, Hestmark G, Ojalora MG, Rauhut A, Büdel B, Scheidegger C, Timdal E, Stenroos S, Brodo I, Perlmutter GB, Ertz D, Diederich P, Lendemer JC, May P, Schoch CL, Arnold AE, Hodkinson BP, Gueidan C, Tripp E, Yahr R, Robertson C, Lutzoni F, 2007 ["2006"]. New insights into classification and evolution of the Lecanoromycetes (Pezizomycotina, Ascomycota) from phylogenetic analyses of three ribosomal RNA- and two protein-coding genes. *Mycologia* 98: 1089–1102.
- Miądlikowska J, Lutzoni F, 2004. Phylogenetic classification of peltigeralean fungi (Peltigerales, Ascomycota) based on ribosomal RNA small and large subunits. *American Journal of Botany* 91: 449–464.
- Miller AN, Huhndorf SM, 2004. A natural classification of Lasiosphaeria based on nuclear LSU rDNA sequences. *Mycological Research* 108: 26–34.
- Miller AN, Huhndorf SM, 2005. Multi-gene phylogenies indicate ascomal wall morphology is a better predictor of phylogenetic relationships than ascospore morphology in the Sordariales (Ascomycota, Fungi). *Molecular Phylogenetics and Evolution* 35: 60–75.
- Miller SL, Larsson E, Larsson K-H, Verbeken A, Nuytinck J, 2007 ["2006"]. Perspectives in the new Russulales. *Mycologia* 98: 960–970.
- Moncalvo J-M, Nilsson RH, Koster B, Dunham SM, Bernauer T, Matheny PB, McLenon T, Margaritescu S, Weiß M, Garnica S, Danell E, Langer G, Langer E, Larsson E, Larsson K-H, Vilgalys R, 2007 ["2006"]. The cantharelloid clade: dealing with incongruent gene trees and phylogenetic reconstruction methods. *Mycologia* 98: 937–948.
- Moncalvo J-M, Vilgalys R, Redhead SA, Johnson JE, James TY, Aime MC, Hofstetter V, Verduin SJW, Larsson E, Baroni TJ, Thorn RG, Jacobsson S, Clémenceçon H, Miller Jr OK, 2002. One hundred seventeen clades of euagarics. *Molecular Phylogenetics and Evolution* 23: 357–400.
- Moore RT, 1980. Taxonomic proposals for the classification of marine yeasts and other yeast-like fungi including the smuts. *Botanica Marina* 23: 361–373.
- Moore RT, 1990. Order Platygloaeales ord. nov. *Mycotaxon* 39: 245–248.
- Moreau F, 1954. *Les Champignons. Physiologie, morphologie, développement et systématique*. Vol. 2. Lechevalier, Paris.
- Morton JB, Benny GL, 1990. Revised classification of arbuscular mycorrhizal fungi (Zygomycetes): a new order, Glomales, two new suborders, Glomineae and Gigasporineae, and two new families, Acaulosporaceae and Gigasporaceae, with an emendation of Glomaceae. *Mycotaxon* 37: 471–491.
- Moss ST, 1975. Septal structure in the Trichomycetes with special reference to *Astreptonema gammari* (Eccriniales). *Transactions of the British Mycological Society* 65: 115–127.
- Nagahama T, Hamamoto M, Nakase T, Shimamura S, Horikoshi K, 2006. Phylogenetic relationship within the Erythrobasidium clade: molecular phylogenies, secondary structure, and intron positions inferred from partial sequences of ribosomal RNA and elongation factor-1 α genes. *Journal of General and Applied Microbiology* 52: 37–45.
- Nagahama T, Sato H, Shimazu M, Sugiyama J, 1995. Phylogenetic divergence of the entomophthoralean fungi: evidence from nuclear 18S ribosomal RNA gene sequences. *Mycologia* 87: 203–209.

- Nannfeldt JA, 1932. Studien über die Morphologie und Systematik der nicht-lichenisierten inoperculaten Discomyceten. *Nova Acta Regiae Societatis Scientiarum Upsaliensis, ser. 4*, 8 (2): 1–368.
- Nishida H, Sugiyama J, 1994. Archiascomycetes: detection of a major new lineage within the Ascomycota. *Mycoscience* 35: 361–366.
- Oberwinkler F, 1976. Eine agaricoide Gattung der *Thelephorales*. *Sydowia* 78: 359–361.
- Oberwinkler F, 1977. Das neue System der Basidiomyceten. In: Frey W, Hurka H, Oberwinkler F (eds), *Beiträge zur Biologie der niederen Pflanzen*. Gustav Fischer Verlag, Stuttgart, pp. 59–105.
- Oberwinkler F, Bandoni RJ, 1982. A taxonomic survey of the gasteroid, auricularioid *Heterobasidiomycetes*. *Canadian Journal of Botany* 60: 1726–1750.
- Oberwinkler F, Bauer R, 1989. The systematics of gasteroid, auricularioid *Heterobasidiomycetes*. *Sydowia* 41: 224–256.
- Oberwinkler F, Bauer R, 1990. *Cryptomycolax*: a new mycoparasitic heterobasidiomycete. *Mycologia* 82: 671–692.
- O'Donnell K, Cigelnik E, Benny GL, 1998. Phylogenetic relationships among the *Harpellales* and *Kickxellales*. *Mycologia* 90: 286–297.
- Pang KL, Abdel-Wahab MA, Sivichai S, El-Sharouney HM, Jones EBG, 2002. *Jahmulales* (*Dothideomycetes*, *Ascomycota*): a new order of lignicolous freshwater ascomycetes. *Mycological Research* 106: 1031–1042.
- Parmasto E, 1986. On the origin of the hymenomycetes (what are corticioid fungi?). *Windahlia* 16: 3–20.
- Peyretailade E, Broussolle V, Peyret P, Méténier G, Gouy M, Vivarès CP, 1998. *Microsporidia*, amitochondrial protists, possess a 70-kDa heat shock protein gene of mitochondrial evolutionary origin. *Molecular Biology and Evolution* 15: 683–689.
- Réblová M, 2006. Molecular systematics of *Ceratostomella sensu lato* and morphologically similar fungi. *Mycologia* 98: 68–93.
- Réblová M, Mostert L, Gams W, Crous PW, 2004. New genera in the *Calosphaerales*: *Togniniella* and its anamorph *Phaeocrella*, and *Calosphaeriophora* as anamorph of *Calosphaeria*. *Studies in Mycology* 50: 533–550.
- Réblová M, Seifert KA, 2004. *Cryptadelphia* (*Trichosphaerales*), a new genus for holomorphs with *Brachysporium* anamorphs and clarification of the taxonomic status of *Wallrothiella*. *Mycologia* 96: 343–367.
- Reeb V, Lutzoni F, Roux C, 2004. Multilocus phylogenetic circumscription of the lichen-forming fungi family *Acarosporaceae* and its position within the *Ascomycota*. *Molecular Phylogenetics and Evolution* 32: 1036–1060.
- Robbertse B, Reeves J, Schoch C, Spatafora JW, 2006. A phylogenomic analysis of the *Ascomycota*. *Fungal Genetics and Biology* 43: 715–725.
- Rossmann AY, Aime MC, Farr DF, Castelbury LA, Peterson KR, Leahy R, 2004. The coelomycetous genera *Chaetomella* and *Pilidium* represent a newly discovered lineage of inoperculate discomycetes. *Mycological Progress* 3: 275–290.
- Ryvarden L, 2002. A note on the genus *Hydnodon* Banker. *Synopsis Fungorum* 15: 31–33.
- Saenz GS, Taylor JW, 1999. Phylogenetic relationships of *Meliola* and *Meliolina* inferred from nuclear small subunit rRNA sequences. *Mycological Research* 103: 1049–1056.
- Saikawa M, Sugiura K, Sato H, 1997. Electron microscopy of two trichomycetous fungi attached to the hindgut lining of pillbugs. *Canadian Journal of Botany* 75: 1479–1484.
- Sampaio JP, Gadanho M, Bauer R, Weiß M, 2003. Taxonomic studies in the *Microbotryomycetidae*: *Leucosporidium golubevii* sp. nov., *Leucosporidiella* gen. nov. and the new orders *Leucosporidiales* and *Sporidiobolales*. *Mycological Progress* 2: 53–68.
- Sampaio JP, 2004. Diversity, phylogeny and classification of basidiomycetous yeasts. In: Agerer R, Piepenbring M, Blanz P (eds), *Frontiers in Basidiomycete Mycology*. IHW Verlag, Eching, pp. 49–80.
- Schaffner JH, 1909. The classification of plants, IV. *Ohio Naturalist* 9: 446–455.
- Schmitt I, Mueller G, Lumbsch HT, 2005. *Ascoma* morphology is homoplaseous and phylogenetically misleading in some pyrenocarpous lichens. *Mycologia* 97: 362–374.
- Schoch CL, Shoemaker RA, Seifert KA, Hambleton S, Spatafora JW, Crous PW, 2007 ['2006']. A multigene phylogeny of the *Dothideomycetes* using four nuclear loci. *Mycologia* 98: 1042–1053.
- Schröter J, 1889. Pilze. In: Cohn F (ed), *Kryptogamen-Flora von Schlesien*, Vol. 3. J. U. Kern's Verlag, Breslau, pp. 382–386.
- Schröter J, 1892. *Phycomycetes*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 63–87.
- Schröter J, 1893. *Monoblepharidaceae*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 106–107.
- Schröter J, 1894. *Pezizineae*. In: Engler A, Prantl K (eds), *Die natürlichen Pflanzenfamilien*, Vol. 1. Wilhelm Engelmann Verlag, Leipzig, pp. 173–243.
- Schüßler A, Schwarzott D, Walker C, 2001. A new fungal phylum, the *Glomeromycota*: phylogeny and evolution. *Mycological Research* 105: 1413–1421.
- Seif E, Leigh L, Roewer I, Forget L, Lang BF, 2005. Comparative mitochondrial genomics in zygomycetes: bacteria-like rRNase P RNAs, mobile elements and a close source of the group I intron invasion in angiosperms. *Nucleic Acids Research* 33: 734–744.
- Serbinow JL, 1907. Organisations I rasiite nekotorich gribow *Chytridinae* Schröt. *Scripa Botanica Horti Universitatis Imperialis Petropolitana* 24: 1–147.
- Shenoy BD, Jeewon R, Wu WP, Bhat DJ, Hyde KD, 2006. Ribosomal and RPB2 DNA sequence analyses suggest that *Sporidesmium* and morphologically similar genera are polyphyletic. *Mycological Research* 110: 916–928.
- Sparrow FK, 1943. *Aquatic Phycomycetes exclusive of the Saprolegniaceae and Pythium*. University of Michigan Press, Ann Arbor.
- Sparrow FK, 1958. Interrelationships and phylogeny of the aquatic *Phycomycetes*. *Mycologia* 50: 797–813.
- Spatafora JW, Johnson D, Sung G-H, Hosaka K, O'Rourke B, Serdani M, Spotts R, Lutzoni F, Hofstetter V, Fraker E, Gueidan C, Miądlikowska J, Reeb V, Lumbsch T, Lücking R, Schmitt I, Aptroot A, Roux C, Miller A, Geiser D, Hafellner J, Hestmark G, Arnold AE, Büdel B, Rauhut A, Hewitt D, Untereiner W, Cole MS, Scheidegger C, Schultz M, Sipman H, Schoch CL, 2007 ['2006']. A five-gene phylogenetic analysis of the *Pezizomycotina*. *Mycologia* 98: 1020–1030.
- Sprague V, Becnel JJ, 1998. Note on the name-author-date combination for the taxon *Microsporidies* Balbiani, 1882, when ranked as a phylum. *Journal of Invertebrate Pathology* 71: 91–94.
- Starbäck K, 1899. *Ascomyceten der Ersten Regnellischen Expedition*. *Bihang till Kungliga Svenska Vetenskaps-Akademiens Handlingar, ser. 3* 25: 1–68.
- Steenkamp ET, Wright J, Baldauf SL, 2006. The protistan origins of animals and fungi. *Molecular Biology and Evolution* 23: 93–106.
- Sugiyama J, Hosaka K, Suh S-O, 2007 ['2006']. Early diverging *Ascomycota*: phylogenetic divergence and related evolutionary enigmas. *Mycologia* 98: 998–1007.
- Suh S-O, Blackwell M, Kurtzman CP, Lachance M-A, 2007 ['2006']. Phylogenetics of *Saccharomycetales*, the ascomycete yeasts. *Mycologia* 98: 1008–1019.
- Swann EC, Frieders EM, McLaughlin DJ, 2001. *Urediniomycetes*. In: McLaughlin DJ, McLaughlin EJ, Lemke P (eds), *The Mycota*. Vol. VII. Part B, *Systematics and Evolution*. Springer-Verlag, Berlin, pp. 37–56.
- Swann EC, Taylor JW, 1995. Phylogenetic perspectives on basidiomycete systematics: evidence from the 18S rRNA gene. *Canadian Journal of Botany* 73: S862–S868.
- Takamatsu S, 2004. Phylogeny and evolution of the powdery mildew fungi (*Erysiphales*, *Ascomycota*) inferred from nuclear ribosomal DNA sequences. *Mycoscience* 45: 147–157.

- Tanabe Y, O'Donnell K, Saikawa M, Sugiyama J, 2000. Molecular phylogeny of parasitic Zygomycota (*Dimargaritales*, *Zoopagales*) based on nuclear small subunit ribosomal DNA sequences. *Molecular Phylogenetics and Evolution* 16: 253–262.
- Tanabe Y, Saikawa M, Watanabe MM, Sugiyama J, 2004. Molecular phylogeny of Zygomycota based on EF-1 and RPB1 sequences: limitations and utility of alternative markers to rDNA. *Molecular Phylogenetics and Evolution* 30: 438–449.
- Tanabe Y, Watanabe MM, Sugiyama J, 2005. Evolutionary relationships among basal fungi (*Chytridiomycota* and *Zygomycota*): Insights from molecular phylogenetics. *Journal of General and Applied Microbiology* 51: 267–276.
- Tang AMC, Jeewon R, Hyde KD, 2007. Phylogenetic utility of protein (RPB2, B-tubulin) and ribosomal (LSU, SSU) gene sequences in the systematics of *Sordariomycetes* (Ascomycota, Fungi). *Antonie van Leeuwenhoek* in press.
- Taylor JW, Spatafora J, O'Donnell K, Lutzoni F, James T, Hibbett DS, Geiser D, Bruns TD, Blackwell M, 2004. The fungi. In: Cracraft J, Donoghue MJ (eds), *Assembling the Tree of Life*. Oxford University Press, Oxford, pp. 171–196.
- Tehler A, 1988. A cladistic outline of the *Eumycota*. *Cladistics* 4: 227–277.
- Tehler A, Little DP, Farris JS, 2003. The full-length phylogenetic tree from 1551 ribosomal sequences of chitinous fungi, *Fungi*. *Mycological Research* 107: 901–916.
- Thorn RG, Moncalvo J-M, Reddy CA, Vilgalys R, 2000. Phylogenetic analyses and the distribution of nematophagy support a monophyletic *Pleurotaceae* within the polyphyletic pleurotoid-lentinoid fungi. *Mycologia* 92: 241–252.
- Tibell L, Vinuesa M, 2005. *Chaenothecopsis* in a molecular phylogeny based on nuclear rDNA ITS and LSU sequences. *Taxon* 54: 427–442.
- Tibell L, Wedin M, 2000. *Mycocaliciales*, a new order for nonlichenized calicioid fungi. *Mycologia* 92: 577–581.
- Underwood LM, 1899. *Moulds, mildews and mushrooms: a guide to the systematic study of the Fungi and Mycetozoa and their literature*. Henry Holt, New York.
- Vijaykrishna D, Mostert L, Jeewon R, Gams W, Hyde KD, Crous PW, 2004. *Pleurostomophora*, an anamorph of *Pleurostoma* (*Calosphaeriales*), a new anamorph genus morphologically similar to *Phialophora*. *Studies in Mycology* 50: 387–395.
- Vossbrinck CR, Debrunner-Vossbrinck BA, 2005. Molecular phylogeny of the *Microsporidia*: ecological, ultrastructural and taxonomic considerations. *Folia Parasitologica* 52: 131–142.
- Wagner T, Fischer M, 2002. Proceedings towards a natural classification of the worldwide taxa *Phellinus* s.l. and *Inonotus* s.l., and phylogenetic relationships of allied genera. *Mycologia* 94: 998–1016.
- Walker C, Schüßler A, 2004. Nomenclatural clarifications and new taxa in the *Glomeromycota*. *Mycological Research* 108: 981–982.
- Wang Z, Binder M, Schoch C, Johnston PR, Spatafora JW, Hibbett DS, 2006. Evolution of helotialean fungi (*Leotiomyces*, *Pezizomycotina*): a nuclear rDNA phylogeny. *Molecular Phylogenetics and Evolution* 41: 295–312.
- Wang Z, Johnston PR, Takamatsu S, Spatafora JW, Hibbett DS, 2007 [‘2006’]. Phylogenetic classification of the *Leotiomyces* based on rDNA data. *Mycologia* 98: 1066–1076.
- Wedin M, Wiklund E, Crewe A, Döring H, Ekman S, Nyberg Å Schmitt I, Lumbsch HT, 2005. Phylogenetic relationships of *Lecanoromycetes* (Ascomycota) as revealed by analyses of mtSSU and nLSU rDNA sequence data. *Mycological Research* 109: 159–172.
- Weir A, Blackwell M, 2001. Molecular data support the *Laboulbeniales* as a separate class of Ascomycota, *Laboulbeniomycetes*. *Mycological Research* 105: 1182–1190.
- Weiß M, Bauer R, Begerow D, 2004a. Spotlights on heterobasidiomycetes. In: Agerer R, Piepenbring M, Blanz P (eds), *Frontiers in Basidiomycote Mycology*. IHW Verlag, Eching, pp. 7–48.
- Weiß M, Oberwinkler F, 2001. Phylogenetic relationships in *Auriculariales* and related groups—hypotheses derived from nuclear ribosomal DNA sequences. *Mycological Research* 105: 403–415.
- Weiß M, Selosse M-A, Rexer K-H, Urban A, Oberwinkler O, 2004b. *Sebacinales*: a hitherto overlooked cosm of heterobasidiomycetes with a broad mycorrhizal potential. *Mycological Research* 108: 1003–1010.
- Wells K, Bandoni RJ, 2001. *Heterobasidiomycetes*. In: McLaughlin DJ, McLaughlin EJ, Lemke P (eds), *The Mycota*. Vol. VII. Part B, *Systematics and Evolution*. Springer-Verlag, Berlin, pp. 85–120.
- White TJ, Bruns TD, Lee S, Taylor JW, 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky J, White TJ (eds), *PCR Protocols: a Guide to Methods and Applications*. Academic Press, San Diego, pp. 315–322.
- White MM, James TY, O'Donnell K, Cafaro MJ, Tanabe Y, Sugiyama J, 2007 [‘2006’]. Phylogeny of the *Zygomycota* based on nuclear ribosomal sequence data. *Mycologia* 98: 872–884.
- Whittaker RH, 1959. On the broad classification of organisms. *Quarterly Review of Biology* 34: 210–226.
- Wiklund E, Wedin M, 2003. The phylogenetic relationships of the cyanobacterial lichens in the *Lecanorales* suborder *Peltigerineae*. *Cladistics* 19: 419–431.
- Wingfield B, Viljoen CD, Wingfield MJ, 1999. Phylogenetic relationships of ophiostomatoid fungi associated with *Protea* infestations in South Africa. *Mycological Research* 103: 1616–1620.
- Winka K, 2000. Phylogenetic relationships within the Ascomycota based on 18S rDNA sequences. PhD thesis, Umeå University, Umeå.
- Winter G, 1880. *Schizomyceten, Saccharomyceten und Basidiomyceten*. In: Rabenhorst's *Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz*, Vol. 1. E. Kummer, Leipzig, pp. 1–924.
- Woronichin NN, 1925. Über die *Capnodiales*. *Annales Mycologici* 23: 174–178.
- Zalar P, de Hoog GS, Schroers H-J, Frank JM, Gunde-Cimerman N, 2005. Taxonomy and phylogeny of the xerophilic genus *Wallemia* (*Wallemiomycetes* and *Wallemiales*, cl. et ord. nov.). *Antonie van Leeuwenhoek* 87: 311–328.
- Zhang N, Castlebury LA, Miller AN, Huhndorf S, Schoch CL, Seifert K, Rossman AY, Rogers JD, Kohlmeyer J, Volkmann-Kohlmeyer B, Sung G-H, 2007 [‘2006’]. *Sordariomycetes* systematics: an overview of the systematics of the *Sordariomycetes* based on a four-gene phylogeny. *Mycologia* 98: 1077–1088.